arXiv:math/0104259v2 [math.NT] 30 Apr 2001

Product formulas on a unitary group in three variables

LEI YANG

1. Introduction

In the present paper, we consider the eigenvalue problems which concern a differential equation

$$L_k f(z_1,z_2)=\lambda f(z_1,z_2). \tag{1.1}$$

Here f is a function on the Siegel domain of type II

$$\mathfrak{S}_2=\{(z_1,z_2)\in\mathbb{C}^2 : z_1+\overline{z_1}-z_2\overline{z_2}>0\}, \tag{1.2}$$

and L_k is a differential operator given by

$$\begin{aligned}L_k&=(z_1+\overline{z_1}-z_2\overline{z_2})\\ &\quad\times\left[(z_1+\overline{z_1})\frac{\partial^2}{\partial z_1\partial\overline{z_1}}+\frac{\partial^2}{\partial z_2\partial\overline{z_2}}+z_2\frac{\partial^2}{\partial\overline{z_1}\partial z_2}+\overline{z_2}\frac{\partial^2}{\partial z_1\partial\overline{z_2}}-k(\frac{\partial}{\partial z_1}-\frac{\partial}{\partial\overline{z_1}})\right],\end{aligned} \tag{1.3}$$

where k is an integer. Let $\Gamma\subset U(2,1)$ be an arithmetic subgroup or a convex cocompact subgroup, where

$$U(2,1)=\{g\in GL(3,\mathbb{C}) : g^*Jg=J\},\quad J=\begin{pmatrix} & & -1\\ & 1 & \\ -1 & & \end{pmatrix}, \tag{1.4}$$

and assume that f is a Γ-automorphic form of weight k in the sense that it is invariant under

$$f\mapsto f(\gamma(z_1,z_2))(a_1z_1+a_2z_2+a_3)^{-k}\overline{(a_1z_1+a_2z_2+a_3)}^{k} \tag{1.5}$$

for every $\gamma=\begin{pmatrix} * & * & *\\ * & * & *\\ a_1 & a_2 & a_3\end{pmatrix}\in\Gamma$. In fact, L_k commutes with (1.5).

Now, let us recall some basic facts about complex hyperbolic geometry (see [Ap] and [Go]). Geometry of complex hyperbolic space $\mathbf{H}^n_{\mathbb{C}}$ is the geometry of the unit ball $\mathbb{B}^n_{\mathbb{C}}$

Typeset by $\mathcal{AMS}$-TEX

in $\mathbb{C}^n$ with the Kähler structure given by the Bergman metric whose automorphisms are biholomorphic automorphisms of the ball, i.e., elements of $PU(n,1)$. Any complex hyperbolic manifold can be represented as the quotient $M = \mathbf{H}^n_{\mathbb{C}}/\Gamma$ by a discrete torsion free isometric action of the fundamental group of M, $\pi_1(M) \cong \Gamma \subset PU(n,1)$, its boundary at infinity $\partial_\infty M$ is naturally identified as the quotient $\Omega(\Gamma)/\Gamma$ of the discontinuity set of Γ at infinity. Here the discontinuity set $\Omega(\Gamma)$ is the maximal subset of $\partial\mathbf{H}^n_{\mathbb{C}}$ where Γ acts discretely, its complement $\Lambda(\Gamma) = \partial\mathbf{H}^n_{\mathbb{C}}\backslash\Omega(\Gamma)$ is the limit set of Γ, $\Lambda(\Gamma) = \overline{\Gamma(x)} \cap \partial\mathbf{H}^n_{\mathbb{C}}$ for any $x \in \mathbf{H}^n_{\mathbb{C}}$.

In general, let G be a connected, linear, real simple Lie group of rank one, $G = KAN$ be an Iwasawa decomposition of G, $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}$ be the corresponding Iwasawa decomposition of the Lie algebra $\mathfrak{g}$, and $P = MAN$ be a minimal parabolic subgroup. The group G acts isometrically on the rank-one symmetric space $X = G/K$. Let $\partial X = G/P = K/M$ be its geodesic boundary. We regard $\overline{X} := X \cup \partial X$ as a compact manifold with boundary.

By the classification of symmetric spaces with strictly negative sectional curvature, we know that X is one of the following spaces: a real hyperbolic space $\mathbf{H}^n_{\mathbb{R}}$ $(n \geq 1)$, a complex hyperbolic space $\mathbf{H}^n_{\mathbb{C}}$ $(n \geq 2)$, a quaternionic hyperbolic space $\mathbf{H}^n_{\mathbb{H}}$ $(n \geq 2)$ or the Cayley hyperbolic plane $\mathbf{H}^2_{\mathbb{O}}$, and G is a linear group finitely covering the orientation-preserving isometric group of X.

Following [BO], we consider a torsion-free discrete subgroup $\Gamma \subset G$ such that ∂X admits a Γ-invariant partition $\partial X = \Omega(\Gamma) \cup \Lambda(\Gamma)$, where $\Omega(\Gamma) \neq \emptyset$ is open and Γ acts freely and cocompactly on $X \cup \Omega(\Gamma)$. The closed subset $\Lambda(\Gamma)$ is called the limit set of Γ. The locally symmetric space $Y := \Gamma\backslash X$ is a complete Riemannian manifold of infinite volume without cusps. It can be compactified by adjoining the geodesic boundary $B := \Gamma\backslash\Omega(\Gamma)$.

A subgroup Γ satisfying this assumption is called convex cocompact or geometrically cocompact since it acts cocompactly on the convex hull of the limit set. The quotient Y is called a Kleinian manifold.

For $H \in \mathfrak{a}$, we define $\rho \in \mathfrak{a}^*$ by $\rho(H) = \frac{1}{2}\mathrm{tr}(\mathrm{ad}(H)_{|\mathfrak{n}})$. Then $\rho(\mathbf{H}^n_{\mathbb{R}}) = \frac{n-1}{2}$ and $\rho(\mathbf{H}^n_{\mathbb{C}}) = n$.

Now, we need the following definition (see [Co]):

Definition. For any discrete group $\Gamma \subset G$, the critical exponent $\delta(\Gamma)$ is the infimum of all s such that

$$\sum_{\gamma\in\Gamma} \exp(-sd(x, \gamma y))$$

converges for some (or any) $x, y \in X$, where $d(x, y)$ is the Riemannian distance from x to y.

The critical exponent $\delta(\Gamma)$ has been extensively studied (see [Pa1], [Su], [Co] and [CoI]). It is known that $\delta(\Gamma) \in [0, 2\rho]$ if Γ is nontrivial. In fact, if $\Omega(\Gamma) \neq \emptyset$, then $\delta(\Gamma)$ is equal to the Hausdorff dimension of the limit set with respect to the natural class of sub-Riemannian metrics on ∂X.

In the case of $U(2,1)$, $\delta(\Gamma) \in [0,4]$ for $X = \mathbf{H}^2_{\mathbb{C}}$.

The theory of Eisenstein series and the spectral theory for Kleinian groups on real hyperbolic spaces $\mathbf{H}^n_{\mathbb{R}}$ has been extensively studied, in particular, by Mandouvalos [Ma1], [Ma2], Patterson [Pa3], and Perry [Pe1], [Pe2], [FHP]. A Kleinian group Γ is a discrete subgroup acting on $\mathbf{H}^n_{\mathbb{R}}$, which acts discontinuously on $\mathbf{H}^n_{\mathbb{R}}$ and has a fundamental domain in $\mathbf{H}^n_{\mathbb{R}}$ of infinite hyperbolic volume. This in particular implies that it is a non-arithmetic subgroup. Their theory extended previous work of Roelcke [Ro], Elstrodt [El], Patterson [Pa2] and Fay [Fa] on the spectral theory and Eisenstein series for Fuchsian groups of the second kind.

Now, we give the corresponding results on the complex hyperbolic space $\mathbf{H}^2_{\mathbb{C}}$. We give the Eisenstein series for the Picard modular group, which is an arithmetic subgroup of $U(2,1)$. We also give the Eisenstein series for any discrete subgroup (either arithmetic or non-arithmetic). In particular, for a convex cocompact subgroup Γ which satisfies that $\delta(\Gamma) < 1$, we give the product formulas. Now, we state the results of this paper.

The Poisson kernel is given by

$$P(Z,W) = \frac{\rho(Z)}{|\rho(Z,W)|^2}, \quad \text{for} \quad Z = (z_1, z_2) \in \mathfrak{S}_2, W = (w_1, w_2) \in \partial\mathfrak{S}_2, \tag{1.6}$$

where

$$\rho(Z,W) := \overline{z_1} + w_1 - \overline{z_2} w_2, \quad \rho(Z) = \rho(Z,Z). \tag{1.7}$$

Let

$$\sigma(Z,Z') := \frac{|\rho(Z,Z')|^2}{\rho(Z)\rho(Z')}. \tag{1.8}$$

It is a point-pair invariant under the action of $U(2,1)$. The critical exponent is given by

$$\delta(\Gamma) := \inf\{s > 0 : \sum_{\gamma\in\Gamma} \sigma(Z, \gamma(Z'))^{-s} < \infty\}. \tag{1.9}$$

For a discrete subgroup $\Gamma \subset U(2,1)$, the Eisenstein series is defined by

$$E(Z,W;s) := \sum_{\gamma\in\Gamma} P(\gamma(Z),W)^s, \quad \text{for} \quad \mathrm{Re}(s) > \delta(\Gamma). \tag{1.10}$$

The automorphic Green function is given by

$$G(Z,Z';s) = \sum_{\gamma\in\Gamma} r(Z,\gamma(Z');s), \quad \text{for} \quad \mathrm{Re}(s) > \delta(\Gamma), \tag{1.11}$$

with

$$r(Z,Z';s)=\frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s-\frac{1}{2})}(\frac{1}{4})^{s-1}u^{-s}F(s,s-1;2s-1;-u^{-1}). \tag{1.12}$$

Here

$$u=u(Z,Z')=\sigma(Z,Z')-1. \tag{1.13}$$

Now, the first main result of this paper is stated as follows:

THEOREM 1.1. *Assume that* Γ *is convex cocompact and* $\delta(\Gamma)<1$. *Then the following product formula holds:*

$$\begin{aligned}&\int_{\Gamma\backslash\Omega(\Gamma)}E(Z,W;s)E(Z',W;2-s)dm(W)\\ &=-\frac{\pi}{s-1}c(s)c(2-s)[G(Z,Z';s)-G(Z,Z';2-s)],\end{aligned} \tag{1.14}$$

where $\delta(\Gamma)<Re(s)<2-\delta(\Gamma)$ *and* $c(s)=\sqrt{\pi}\Gamma(s-\frac{1}{2})\Gamma(s)^{-1}$.

For $\gamma=\begin{pmatrix}* & * & *\\ * & * & *\\ a_1 & a_2 & a_3\end{pmatrix}\in U(2,1)$, set

$$j(\gamma,Z)=a_1z_1+a_2z_2+a_3z_3. \tag{1.15}$$

The S-matrix is given by

$$S(W,W';s):=\sum_{\gamma\in\Gamma}|\rho(\gamma(W),W')|^{-2s}|j(\gamma,W)|^{-2s} \tag{1.16}$$

for $W,W'\in\Omega(\Gamma)$ and $\mathrm{Re}(s)>\delta(\Gamma)$. By the S-matrix, we give the functional equation of Eisenstein series:

THEOREM 1.2. *Assume that* Γ *is convex cocompact and* $\delta(\Gamma)<1$. *Then the following functional equation for Eisenstein series holds:*

$$\int_{\Gamma\backslash\Omega(\Gamma)}E(Z,W;s)S(W,W';2-s)dm(W)=\frac{4^{s-1}\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}E(Z,W';2-s), \tag{1.17}$$

for $1<Re(s)<2-\delta(\Gamma)$.

The Poisson kernel of weight k is given by

$$P_k(Z,W;s)=\rho(Z)^s\rho(Z,W)^{k-s}\rho(\overline{Z},\overline{W})^{-k-s}, \tag{1.18}$$

where $Z \in \mathfrak{S}_2$, $W \in \partial\mathfrak{S}_2$. The Eisenstein series of weight k is defined by

$$E(Z,W;k,s)=\sum_{\gamma\in\Gamma} j(\gamma,Z)^{-k} j(\overline{\gamma},\overline{Z})^{k} P_k(\gamma(Z),W;s), \quad \text{for} \quad \mathrm{Re}(s)>\delta(\Gamma). \tag{1.19}$$

Set

$$\begin{aligned} K(Z,W;k,s)=&\frac{\pi^{\frac{3}{2}}\Gamma(\frac{3}{2}-s)}{(|k|+1-s)\Gamma(2-s)}(\frac{1}{4})^{s-1}\rho(Z,W)^{k}\rho(\overline{Z},\overline{W})^{-k}\\ &\times \sigma^{-|k|}(\sigma-1)^{|k|-s}F(s-|k|,s-1-|k|;2s-1;-\frac{1}{\sigma-1}). \end{aligned} \tag{1.20}$$

The automorphic Green function of weight k is given by

$$G(Z,Z';k,s)=\sum_{\gamma\in\Gamma} K(Z,\gamma(Z');k,s), \quad \text{for} \quad \mathrm{Re}(s)>\delta(\Gamma). \tag{1.21}$$

Now, we state the main theorem of this paper:

THEOREM 1.3 (MAIN THEOREM). *Assume that* Γ *is convex cocompact and* $\delta(\Gamma)<1$. *Then the following product formula holds:*

$$\begin{aligned} &\int_{\Gamma\backslash\Omega(\Gamma)} E(Z,W;k,s)E(Z',W;-k,2-s)dm(W)\\ =&G(Z,Z';k,s)+G(Z,Z';k,2-s)\\ =&G(Z',Z;-k,s)+G(Z',Z;-k,2-s), \end{aligned} \tag{1.22}$$

for $\delta(\Gamma)<Re(s)<2-\delta(\Gamma)$ *and* $k=0$ *or* $k=\pm 1$.

The S-matrix of weight k is defined by

$$S(W,W';k,s):=\sum_{\gamma\in\Gamma} j(\gamma,W)^{k-s} j(\overline{\gamma},\overline{W})^{-k-s}\rho(\gamma(W),W')^{-k-s}\rho(\overline{\gamma(W)},\overline{W'})^{k-s} \tag{1.23}$$

for $W,W'\in\Omega(\Gamma)$ and $\mathrm{Re}(s)>\delta(\Gamma)$. By the S-matrix of weight k, we obtain the functional equation of Eisenstein series of weight k.

THEOREM 1.4. *Assume that* Γ *is convex cocompact and* $\delta(\Gamma)<1$. *Then the following functional equation for Eisenstein series of weight* k *holds:*

$$\begin{aligned} &\int_{\Gamma\backslash\Omega(\Gamma)} E(Z,W;k,s)S(W,W';-k,2-s)dm(W)\\ =&\frac{\pi^{\frac{3}{2}}\Gamma(s-\frac{1}{2})}{(|k|+s-1)\Gamma(s)}4^{s-1}E(Z,W';k,2-s), \end{aligned} \tag{1.24}$$

for $1 < Re(s) < 2 - \delta(\Gamma)$ *and* $k = 0, \pm 1$.

In fact, when $k = 0$, Theorem 1.3 and Theorem 1.4 reduces to Theorem 1.1 and Theorem 1.2.

In the appendix, we give the product formulas on $SL(2, \mathbb{R})$.

THEOREM 1.5. *Assume that* Γ *is convex cocompact and* $\delta(\Gamma) < \frac{1}{2}$. *Then the following product formula on* $SL(2, \mathbb{R})$ *holds:*

$$\int_{\Gamma\backslash\Omega(\Gamma)} E(z, \zeta; k, s)E(z', \zeta; -k, 1-s)dm(\zeta) = G(z, z'; k, s) + G(z, z'; k, 1-s), \tag{1.25}$$

for $\delta(\Gamma) < Re(s) < 1 - \delta(\Gamma)$ *and* $k = 0, \pm 1$.

2. Epstein zeta function and Eisenstein series on Picard modular groups

Let $\mathfrak{S}_2$ be the Siegel domain

$$\mathfrak{S}_2 = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 + \overline{z_1} > |z_2|^2\}. \tag{2.1}$$

The imaginary quadratic field $K = \mathbb{Q}(\sqrt{-3})$ is called the field of Eisenstein numbers. Its ring of integers

$$\mathcal{O} = \mathcal{O}_K = \{a/2 + b\sqrt{-3}/2 : a, b \in \mathbb{Z}, a \equiv b (\mathrm{mod} 2)\} = \mathbb{Z} + \mathbb{Z}[\omega] \tag{2.2}$$

with $\omega = \frac{-1+\sqrt{-3}}{2}$ is called the ring of Eisenstein integers.

In their paper [KW], Korányi and Wolf studied the general Cayley transform which carries the bounded domain of the Harish-Chandra realization into a generalized half-plane. In [KR], Korányi and Reimann gave the Cayley transform on $U(2, 1)$. Now, we define the Cayley transform $C : \mathbb{B}^2 := \{(w_1, w_2) \in \mathbb{C}^2 : |w_1|^2 + |w_2|^2 < 1\} \to \mathfrak{S}_2$, where

$$C = \begin{pmatrix} -\overline{\omega} & & -\omega \\ & 1 & \\ -1 & & 1 \end{pmatrix} \in GL(3, \mathcal{O}). \tag{2.3}$$

It is known that

$$U(2, 1) = \{g \in GL(3, \mathbb{C}) : gI_{2,1}g^* = I_{2,1}\}, \tag{2.4}$$

where $I_{2,1} = \mathrm{diag}\{1, 1, -1\}$.

Let $E_{i,j}$ be the 3×3 matrix with a 1 at the i, j-th component, zeros elsewhere. Using Kronecker's delta, we have

$$[E_{p,q}, E_{r,s}] = \delta_{q,r}E_{p,s} - \delta_{s,p}E_{r,q}.$$

A basis of $\mathfrak{g} = u(2,1)$ is as follows:

$$X_1 = E_{2,3} + E_{3,2}, \quad X_2 = E_{1,3} + E_{3,1}, \quad X_3 = E_{1,2} - E_{2,1},$$
$$X_4 = iE_{2,3} - iE_{3,2}, \quad X_5 = iE_{1,3} - iE_{3,1}, \quad X_6 = iE_{1,2} + iE_{2,1},$$
$$X_7 = iE_{1,1}, \quad X_8 = iE_{2,2}, \quad X_9 = iE_{3,3}.$$

Thus,

$$E_{2,3} = \frac{1}{2}(X_1 - iX_4), \quad E_{3,2} = \frac{1}{2}(X_1 + iX_4),$$
$$E_{1,3} = \frac{1}{2}(X_2 - iX_5), \quad E_{3,1} = \frac{1}{2}(X_2 + iX_5),$$
$$E_{1,2} = \frac{1}{2}(X_3 - iX_6), \quad E_{2,1} = -\frac{1}{2}(X_3 + iX_6),$$
$$E_{1,1} = -iX_7, \quad E_{2,2} = -iX_8, \quad E_{3,3} = -iX_9.$$

The action of $U(2,1)$ on $\mathbb{B}^2$ is as follows:

$$\begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} (w_1, w_2) = \left(\frac{a_1 w_1 + a_2 w_2 + a_3}{c_1 w_1 + c_2 w_2 + c_3}, \frac{b_1 w_1 + b_2 w_2 + b_3}{c_1 w_1 + c_2 w_2 + c_3} \right).$$

We have

$$\begin{aligned}
&X_3 f(w_1, w_2, \overline{w_1}, \overline{w_2}) \\
=&\frac{d}{dt}[f(\exp(t \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix})(w_1, w_2)) + f(\exp(t \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix})(\overline{w_1}, \overline{w_2}))]|_{t=0} \\
=&\frac{d}{dt}[f(\begin{pmatrix} \cos t & \sin t & 0 \\ -\sin t & \cos t & 0 \\ 0 & 0 & 1 \end{pmatrix} (w_1, w_2)) + f(\begin{pmatrix} \cos t & \sin t & 0 \\ -\sin t & \cos t & 0 \\ 0 & 0 & 1 \end{pmatrix} (\overline{w_1}, \overline{w_2}))]|_{t=0} \\
=&\frac{d}{dt} f(w_1 \cos t + w_2 \sin t, -w_1 \sin t + w_2 \cos t)|_{t=0} \\
&+ \frac{d}{dt} f(\overline{w_1} \cos t + \overline{w_2} \sin t, -\overline{w_1} \sin t + \overline{w_2} \cos t)|_{t=0} \\
=&w_2 \frac{\partial f}{\partial w_1} - w_1 \frac{\partial f}{\partial w_2} + \overline{w_2} \frac{\partial f}{\partial \overline{w_1}} - \overline{w_1} \frac{\partial f}{\partial \overline{w_2}}.
\end{aligned}$$

So, the corresponding operator is

$$X_3 = w_2 \frac{\partial}{\partial w_1} - w_1 \frac{\partial}{\partial w_2} + \overline{w_2} \frac{\partial}{\partial \overline{w_1}} - \overline{w_1} \frac{\partial}{\partial \overline{w_2}}.$$

Similarly, we have

$$X_2 = (1 - w_1^2)\frac{\partial}{\partial w_1} - w_1 w_2 \frac{\partial}{\partial w_2} + (1 - \overline{w_1}^2)\frac{\partial}{\partial \overline{w_1}} - \overline{w_1 w_2}\frac{\partial}{\partial \overline{w_2}}.$$

$$X_1 = -w_1 w_2 \frac{\partial}{\partial w_1} + (1 - w_2^2)\frac{\partial}{\partial w_2} - \overline{w_1 w_2}\frac{\partial}{\partial \overline{w_1}} + (1 - \overline{w_2}^2)\frac{\partial}{\partial \overline{w_2}}.$$

$$X_6 = iw_2 \frac{\partial}{\partial w_1} + iw_1 \frac{\partial}{\partial w_2} - i\overline{w_2}\frac{\partial}{\partial \overline{w_1}} - i\overline{w_1}\frac{\partial}{\partial \overline{w_2}}.$$

$$X_5 = i(1 + w_1^2)\frac{\partial}{\partial w_1} + iw_1 w_2 \frac{\partial}{\partial w_2} - i(1 + \overline{w_1}^2)\frac{\partial}{\partial \overline{w_1}} - i\overline{w_1 w_2}\frac{\partial}{\partial \overline{w_2}}.$$

$$X_4 = iw_1 w_2 \frac{\partial}{\partial w_1} + i(1 + w_2^2)\frac{\partial}{\partial w_2} - i\overline{w_1 w_2}\frac{\partial}{\partial \overline{w_1}} - i(1 + \overline{w_2}^2)\frac{\partial}{\partial \overline{w_2}}.$$

$$X_7 = iw_1 \frac{\partial}{\partial w_1} - i\overline{w_1}\frac{\partial}{\partial \overline{w_1}}.$$

$$X_8 = iw_2 \frac{\partial}{\partial w_2} - i\overline{w_2}\frac{\partial}{\partial \overline{w_2}}.$$

$$X_9 = -iw_1 \frac{\partial}{\partial w_1} - iw_2 \frac{\partial}{\partial w_2} + i\overline{w_1}\frac{\partial}{\partial \overline{w_1}} + i\overline{w_2}\frac{\partial}{\partial \overline{w_2}}.$$

We denote X_{ij} as the differential operator corresponding to $E_{i,j}$, then

$$X_{23} = \frac{1}{2}(X_1 - iX_4) = \frac{\partial}{\partial w_2} - \overline{w_1 w_2}\frac{\partial}{\partial \overline{w_1}} - \overline{w_2}^2 \frac{\partial}{\partial \overline{w_2}}, \quad X_{32} = \frac{1}{2}(X_1 + iX_4) = \overline{X_{23}},$$

$$X_{13} = \frac{1}{2}(X_2 - iX_5) = \frac{\partial}{\partial w_1} - \overline{w_1}^2 \frac{\partial}{\partial \overline{w_1}} - \overline{w_1 w_2}\frac{\partial}{\partial \overline{w_2}}, \quad X_{31} = \frac{1}{2}(X_2 + iX_5) = \overline{X_{13}},$$

$$X_{12} = \frac{1}{2}(X_3 - iX_6) = w_2 \frac{\partial}{\partial w_1} - \overline{w_1}\frac{\partial}{\partial \overline{w_2}}, \quad X_{21} = -\frac{1}{2}(X_3 + iX_6) = -\overline{X_{12}},$$

$$X_{11} = -iX_7 = w_1 \frac{\partial}{\partial w_1} - \overline{w_1}\frac{\partial}{\partial \overline{w_1}},$$

$$X_{22} = -iX_8 = w_2 \frac{\partial}{\partial w_2} - \overline{w_2}\frac{\partial}{\partial \overline{w_2}},$$

$$X_{33} = -iX_9 = -w_1 \frac{\partial}{\partial w_1} - w_2 \frac{\partial}{\partial w_2} + \overline{w_1}\frac{\partial}{\partial \overline{w_1}} + \overline{w_2}\frac{\partial}{\partial \overline{w_2}}.$$

Now, if n is any positive integer, we claim that

$$\sum_{i_1=1}^{3}\sum_{i_2=1}^{3}\cdots\sum_{i_n=1}^{3} X_{i_1 i_2} X_{i_2 i_3} \cdots X_{i_n i_1}$$

lies in the center of $U(\mathfrak{g})$. To prove this, without loss of generality, let us assume $n = 2$. Then

$$[X_{pq}, \sum_{i_1}\sum_{i_2} X_{i_1i_2}X_{i_2i_1}] = \sum_{i_1}\sum_{i_2}([X_{pq}, X_{i_1i_2}]X_{i_2i_1} + X_{i_1i_2}[X_{pq}, X_{i_2i_1}]).$$

By $[X_{pq}, X_{rs}] = \delta_{qr}X_{ps} - \delta_{sp}X_{rq}$, this equals

$$\begin{aligned}
&\sum_{i_1}\sum_{i_2}(\delta_{qi_1}X_{pi_2}X_{i_2i_1} - \delta_{i_2p}X_{i_1q}X_{i_2i_1} + X_{i_1i_2}\delta_{qi_2}X_{pi_1} - X_{i_1i_2}\delta_{i_1p}X_{i_2q})\\
=&\sum_{i_2} X_{pi_2}X_{i_2q} - \sum_{i_1} X_{i_1q}X_{pi_1} + \sum_{i_1} X_{i_1q}X_{pi_1} - \sum_{i_2} X_{pi_2}X_{i_2q} = 0.
\end{aligned}$$

Thus the above sum commutes with the generators X_i of $U(\mathfrak{g})$, and lies in the center of $U(\mathfrak{g})$. The proof is similar if $n \neq 2$.

Set

$$D_1 = \sum_{i_1=1}^{3}\sum_{i_2=1}^{3} E_{i_1i_2}E_{i_2i_1},$$

$$D_2 = \sum_{i_1=1}^{3}\sum_{i_2=1}^{3}\sum_{i_3=1}^{3} E_{i_1i_2}E_{i_2i_3}E_{i_3i_1}.$$

In the category of differential operators,

$$[X_{pq}, X_{rs}] = \delta_{sp}X_{rq} - \delta_{qr}X_{ps}.$$

Therefore, we have

$$D_1 = 2(X_{11}^2 + X_{22}^2 + X_{11}X_{22} + X_{12}X_{21} + X_{23}X_{32} + X_{31}X_{13}) = 2\Delta,$$

$$\begin{aligned}
D_2 =&3[X_{11}(X_{12}X_{21} - X_{32}X_{23} - X_{22}^2) + X_{22}(X_{21}X_{12} - X_{31}X_{13} - X_{11}^2)\\
&+ X_{23}X_{31}X_{12} + X_{13}X_{32}X_{21} - X_{31}X_{13} - X_{32}X_{23} + X_{11} + X_{22}] = -3\Delta,
\end{aligned}$$

where

(2.5)

$$\begin{aligned}
\Delta =& (1 - |w_1|^2 - |w_2|^2)\\
&\times \left[(1 - |w_1|^2)\frac{\partial^2}{\partial w_1 \partial \overline{w_1}} + (1 - |w_2|^2)\frac{\partial^2}{\partial w_2 \partial \overline{w_2}} - w_1\overline{w_2}\frac{\partial^2}{\partial w_1 \partial \overline{w_2}} - \overline{w_1}w_2\frac{\partial^2}{\partial \overline{w_1} \partial w_2}\right]
\end{aligned}$$

is the Laplace-Beltrami operator of $U(2,1)$ on $\mathbb{B}^2$.

Let $G = CU(2,1)C^{-1}$, $K = C(U(2) \times U(1))C^{-1}$ and $G(\mathcal{O}) = CU(2,1;\mathcal{O})C^{-1}$, then $G(\mathcal{O})$ is an arithmetic subgroup of G and $S := G(\mathcal{O})\backslash G/K$ is called a Picard modular surface (see [Pic1], [Pic2], [Gir], [Ho1], [Ho2], [LR] and [BB] for the details).

The complex hyperbolic Gauss-Bonnet formula states that (see [HP] and [P])

$$\mathrm{Vol}(\mathbb{H}^2_{\mathbb{C}}/\Gamma) = \frac{8\pi^2}{3}\chi(\mathbb{H}^2_{\mathbb{C}}/\Gamma),$$

where $\chi(\mathbb{H}^2_{\mathbb{C}}/\Gamma)$ is the Euler number of $\mathbb{H}^2_{\mathbb{C}}/\Gamma$. Holzapfel showed that

$$\chi(\mathbb{B}^2/PSU(2,1;\mathcal{O})) = \frac{1}{24}.$$

As $PSU(2,1;\mathcal{O})$ has index 3 in $PU(2,1;\mathcal{O})$, we see that $\chi(\mathbb{B}^2/PU(2,1;\mathcal{O})) = \frac{1}{72}$, and so

$$\mathrm{Vol}_{\mathbb{B}^2}(\mathbb{B}^2/PU(2,1;\mathcal{O})) = \frac{8\pi^2}{3}\cdot\frac{1}{72} = \frac{\pi^2}{27}.$$

Denote $J = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$, then $CI_{2,1}C^* = J$. It follows that

$$G = \{g \in GL(3,\mathbb{C}) : gJg^* = J\}. \tag{2.6}$$

This is the other realization of $U(2,1)$. For simplicity, from now on, we use the same symbol $U(2,1)$ to denote as G. In fact, $G = \{g \in GL(3,\mathbb{C}) : g^*Jg = J\}$.

Let P be the parabolic subgroup of $U(2,1)$ which consists of upper triangular matrices. This group P has the Langlands decomposition $P = NAM$, where

$$N = \left\{ n = [z,t] = \begin{pmatrix} 1 & z & \frac{1}{2}|z|^2 + it \\ 0 & 1 & \overline{z} \\ 0 & 0 & 1 \end{pmatrix} : z \in \mathbb{C}, t \in \mathbb{R} \right\}.$$

$$A = \left\{ a = \begin{pmatrix} \rho^{-1} & & \\ & 1 & \\ & & \rho \end{pmatrix} : \rho > 0 \right\},$$

$$M = \left\{ m = \begin{pmatrix} \beta & & \\ & \beta^{-2} & \\ & & \beta \end{pmatrix} : \beta \in \mathbb{C}, |\beta|^2 = 1 \right\}.$$

Then $G = NAMK = NAK$ is the Iwasawa decomposition. Set $\Gamma = G(\mathcal{O})$, then

$$\Gamma \cap N = \left\{ [\alpha,\beta] = \begin{pmatrix} 1 & \alpha & \beta \\ & 1 & \overline{\alpha} \\ & & 1 \end{pmatrix} : |\alpha|^2 = \beta + \overline{\beta}, \alpha, \beta \in \mathcal{O} \right\}. \tag{2.7}$$

If $k \in K$, then

$$k = C \begin{pmatrix} \alpha & \beta & 0 \\ \gamma & \delta & 0 \\ 0 & 0 & e^{i\theta} \end{pmatrix} C^{-1},$$

where $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in U(2)$ and $\theta \in [0, 2\pi)$. Write explicitly,

$$k = \begin{pmatrix} -\overline{\omega}\alpha - \omega e^{i\theta} & -\overline{\omega}\beta & -\alpha + e^{i\theta} \\ \gamma & \delta & \omega\gamma \\ -\alpha + e^{i\theta} & -\beta & -\omega\alpha - \overline{\omega}e^{i\theta} \end{pmatrix}.$$

We have $k(-\omega, 0) = (-\omega, 0)$. Note that $\rho(-\omega, 0) = 1$. Therefore, $g(-\omega, 0) = (-\omega, 0)$ if and only if $g \in K$. Define $\pi : G \to \mathfrak{S}_2$ by $\pi(g) = g(-\omega, 0)$. It induces a bijection $G/K \to \mathfrak{S}_2$. For $g = nak$,

$$\pi(g) = nak(-\omega, 0) = na(-\omega, 0) = (-\omega\rho^{-2} + \frac{1}{2}|z|^2 + it, \overline{z}).$$

Thus, if we set $(z_1, z_2) = \pi(g)$, then $z_1 + \overline{z_1} - z_2\overline{z_2} = \rho^{-2}$.

Following [He], let $\Delta = \Delta(\Gamma) = \left\{ \begin{pmatrix} 1 & & ix \\ & 1 & \\ & & 1 \end{pmatrix} : x \in \mathbb{R} \right\} \cap \Gamma$ and $q = q(\Gamma)$ be the positive integer such that $\begin{pmatrix} 1 & & q\sqrt{-3} \\ & 1 & \\ & & 1 \end{pmatrix}$ generates Δ. For $[\alpha, \beta], [\alpha', \beta'] \in \Gamma \cap N$,

$$[\alpha', \beta'][\alpha, \beta]([\alpha, \beta][\alpha', \beta'])^{-1} = [0, 2\sqrt{-3}\mathrm{Im}(\alpha'\overline{\alpha})] \in \Delta,$$

where $\mathrm{Im}\alpha := b$ for $\alpha = a + b\sqrt{-3}$, $a, b \in \mathbb{Z}$. Hence, $2q^{-1}\mathrm{Im}(\alpha'\overline{\alpha}) \in \mathbb{Z}$.

By Cayley transform, we have

THEOREM 2.1. *The Laplace-Beltrami operator of* $U(2,1)$ *on* $\mathfrak{S}_2$ *is*

$$L = (z_1 + \overline{z_1} - |z_2|^2) \left[(z_1 + \overline{z_1}) \frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2} \frac{\partial^2}{\partial z_1 \partial \overline{z_2}} \right]. \tag{2.8}$$

If $g = \begin{pmatrix} 1 & \alpha & \beta \\ & 1 & \gamma \\ & & 1 \end{pmatrix} \in \Gamma$, then $\alpha = \overline{\gamma}$, $|\alpha|^2 = \beta + \overline{\beta}$. Denote $Z = (z_1, z_2)$, set $\rho(Z) = z_1 + \overline{z_1} - |z_2|^2$, we have

$$\rho(g, Z) := \rho(g(Z)) = (z_1 + \alpha z_2 + \beta) + (\overline{z_1} + \overline{\alpha z_2} + \overline{\beta}) - (z_2 + \gamma)(\overline{z_2} + \overline{\gamma}) = \rho(Z).$$

If $g = \begin{pmatrix} * & * & * \\ * & * & * \\ a_1 & a_2 & a_3 \end{pmatrix} \in U(2,1)$, then $a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}$. By a straightforward calculation, we have

$$\rho(g, Z) = \frac{\rho(Z)}{|j(\gamma, Z)|^2}, \quad \text{with} \quad j(\gamma, Z) := a_1z_1 + a_2z_2 + a_3. \tag{2.9}$$

For the Picard modular group $\Gamma = U(2, 1; \mathcal{O})$, the Eisenstein series is defined as

$$E(Z; s) = \sum_{\gamma \in \Gamma \cap N \backslash \Gamma} \rho(\gamma, Z)^s, \quad \text{for} \quad \mathrm{Re}(s) > 2. \tag{2.10}$$

The unit group of $\mathcal{O}$ is $\{\pm 1, \pm\omega, \pm\overline{\omega}\}$, its order is 6.

$$E(Z; s) = \frac{1}{6} \sum_{\substack{a_1, a_2, a_3 \in \mathcal{O}, (a_1, a_2, a_3) = \mathcal{O}, \\ a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}}} \frac{\rho^s}{|a_1z_1 + a_2z_2 + a_3|^{2s}},$$

where (a_1, a_2, a_3) denotes the ideal generated by a_1, a_2 and a_3.

Theorem 2.2. *The Eisenstein series satisfies the following equation:*

$$LE(Z; s) = s(s - 2)E(Z; s). \tag{2.11}$$

Proof. For $j(\gamma, Z) = a_1z_1 + a_2z_2 + a_3$,

$$L\rho(\gamma, Z)^s = s(s - 2)\rho(\gamma, Z)^s + (-\overline{a_1}a_3 - a_1\overline{a_3} + a_2\overline{a_2})s^2 \frac{\rho(Z)^s}{|j(\gamma, Z)|^{2(s+1)}}.$$

□

Let

$$\mathcal{SP}_3(\mathbb{C}) = \{Y \in M_3(\mathbb{C}) : Y \quad \text{positive definite}, \quad Y^* = Y, \det Y = 1\},$$

$\mathcal{H}_3(\mathbb{C}) := \mathcal{SP}_3(\mathbb{C}) \cap U(2, 1)$ and denote $Y[a] = a^*Ya$, for $Y \in \mathcal{SP}_3(\mathbb{C})$.

For $a \in A, n \in N$ as above,

$$a[n] := n^*an = \begin{pmatrix} \frac{1}{\rho} & \frac{z}{\rho} & \frac{\frac{1}{2}|z|^2 + it}{\rho} \\ \frac{\overline{z}}{\rho} & \frac{|z|^2}{\rho} + 1 & \overline{z}(\frac{\frac{1}{2}|z|^2 + it}{\rho} + 1) \\ \frac{\frac{1}{2}|z|^2 - it}{\rho} & z(\frac{\frac{1}{2}|z|^2 - it}{\rho} + 1) & \frac{\frac{1}{4}|z|^4 + t^2}{\rho} + |z|^2 + \rho \end{pmatrix}.$$

We can identify $\mathfrak{S}_2$ with $\mathcal{H}_3(\mathbb{C})$ by $(z_1, z_2) \mapsto W_{(\rho,t,z)} = a[n]$, where

$$z_1 = \rho + \frac{1}{2}|z|^2 - it, z_2 = -\overline{z}, \rho = \frac{1}{2}(z_1 + \overline{z_1} - |z_2|^2).$$

LEMMA 2.3. *If* $a = (a_3, -a_2, a_1) \in \mathcal{O}^3$, *where* $a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}$, *then*

$$W_{(\rho,t,z)}[a^*] = \rho^{-1}|a_1 z_1 + a_2 z_2 + a_3|^2. \tag{2.12}$$

Proof. In fact,

$$\rho^{-1}|a_1 z_1 + a_2 z_2 + a_3|^2 - W_{(\rho,t,z)}[a^*] = a_1\overline{a_3} + \overline{a_1}a_3 - a_2\overline{a_2} = 0.$$

□

Definition 2.4. For the Picard modular group $\Gamma = U(2,1;\mathcal{O})$ the Epstein zeta function is given by

$$Z(Y, s) := \frac{1}{6} \sum_{a \in \mathcal{O}^3 - \{0\}, a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}} Y[a]^{-s}, \tag{2.13}$$

where $\mathrm{Re}(s) > 2$.

LEMMA 2.5. $Z(W_{(\rho,t,z)}, s) = \zeta_K(2s)E(z_1, z_2; s)$, *where* $\zeta_K(s)$ *is the Dedekind zeta function of the number field* $K = \mathbb{Q}(\sqrt{-3})$.

Because $E(z_1, z_2; s) = Z(W_{(\rho,t,z)}, s)/\zeta_K(2s)$, any zero of $\zeta_K(2s)$ will give a pole of $E(z_1, z_2; s)$ unless $Z(W, s)$ vanishes at that value of s. The trivial zeros of $Z(W, s)$ and $\zeta_K(2s)$ are both $s = -1, -2, -3, \cdots$, and both have order 1. So the trivial zeros cancel out in $E(z_1, z_2; s)$.

Suppose $a_1, a_2, a_3 \in \mathcal{O}$ and $\overline{a_1}a_3 + a_1\overline{a_3} = a_2\overline{a_2}$, then $J[(a_1, a_2, a_3)^*] = 0$ and $J[(a_3, -a_2, a_1)^*] = 0$.

For n, $a \in U(2,1)$ as above, then $n^* \in U(2,1)$, thus $W \in U(2,1)$. If $g \in \mathcal{H}_3(\mathbb{C})$, $gJg = J$, thus $g^{-1} = JgJ$. Suppose $a = (a_1, a_2, a_3) \in \mathcal{O}^3 - \{0\}$, we have the following lemma:

LEMMA 2.6. *For* $g \in \mathcal{H}_3(\mathbb{C})$, $g[(a_3, -a_2, a_1)^*] = g^{-1}[(a_1, a_2, a_3)^*]$.

Proof. In fact,

$$g^{-1}[(a_1, a_2, a_3)^*] = JgJ[(a_1, a_2, a_3)^*] = (-a_3, a_2, -a_1)g \begin{pmatrix} -\overline{a_3} \\ \overline{a_2} \\ -\overline{a_1} \end{pmatrix} = g[(a_3, -a_2, a_1)^*].$$

□

By Lemma 2.6, it follows that

Theorem 2.7. *The functional equation of Epstein zeta functions:*

$$Z(Y^{-1}, s) = Z(Y, s) \tag{2.14}$$

if $Y \in \mathcal{H}_3(\mathbb{C})$.

Following [Ho2], we define the Picard modular forms on $\mathfrak{S}_2$ as follows:

Definition 2.8. A holomorphic function $f : \mathfrak{S}_2 \to \mathbb{C}$ is a Picard modular form of $K = \mathbb{Q}(\sqrt{-3})$ and of weight k, if it satisfies:

$$f(\gamma(Z)) = \mathrm{Jac}(\gamma, Z)^{-k} f(Z) \quad \text{for all} \quad \gamma \in \Gamma, \tag{2.15}$$

where $\mathrm{Jac}(\gamma, Z)$ is the Jacobi determinant of $\gamma : \mathfrak{S}_2 \to \mathfrak{S}_2$ at $Z = (z_1, z_2)$.

Now, for the Picard modular group $\Gamma = U(2, 1; \mathcal{O})$, we define the Eisenstein series of weight k:

$$E_k(Z) = \sum_{\gamma \in \Gamma \cap N \backslash \Gamma} \mathrm{Jac}(\gamma, Z)^k, \quad \text{for} \quad k > 1. \tag{2.16}$$

In fact, in [BB] and [Sh1], the authors studied the Poincaré-Eisenstein series.

The following proposition is important.

Proposition 2.9. *$j(g, Z)$ satisfies the following identity:*

$$j(g_1 g_2, Z) = j(g_1, g_2(Z)) j(g_2, Z). \tag{2.17}$$

Proof. For $g_1 = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in U(2, 1)$ and $g_2 = \begin{pmatrix} a'_1 & a'_2 & a'_3 \\ b'_1 & b'_2 & b'_3 \\ c'_1 & c'_2 & c'_3 \end{pmatrix} \in U(2, 1)$, we have

$$g_1 g_2 = \begin{pmatrix} * & * & * \\ * & * & * \\ a'_1 c_1 + b'_1 c_2 + c'_1 c_3 & a'_2 c_1 + b'_2 c_2 + c'_2 c_3 & a'_3 c_1 + b'_3 c_2 + c'_3 c_3 \end{pmatrix}.$$

Thus,

$$j(g_1 g_2, Z) = (a'_1 c_1 + b'_1 c_2 + c'_1 c_3) z_1 + (a'_2 c_1 + b'_2 c_2 + c'_2 c_3) z_2 + (a'_3 c_1 + b'_3 c_2 + c'_3 c_3).$$

On the other hand.

$$g_2(Z) = \left(\frac{a'_1 z_1 + a'_2 z_2 + a'_3}{c'_1 z_1 + c'_2 z_2 + c'_3}, \frac{b'_1 z_1 + b'_2 z_2 + b'_3}{c'_1 z_1 + c'_2 z_2 + c'_3} \right).$$

$$j(g_1, g_2(Z)) = c_1 \frac{a_1' z_1 + a_2' z_2 + a_3'}{c_1' z_1 + c_2' z_2 + c_3'} + c_2 \frac{b_1' z_1 + b_2' z_2 + b_3'}{c_1' z_1 + c_2' z_2 + c_3'} + c_3.$$

$$j(g_2, Z) = c_1' z_1 + c_2' z_2 + c_3'.$$

So,

$$j(g_1, g_2(Z)) j(g_2, Z) = j(g_1 g_2, Z).$$

□

3. Product formulas on $U(2,1)$

In his paper [Ko], the Poisson kernel $P : B \times D \to \mathbb{R}$ of D is defined by

$$P(u, z) = \frac{|S(u, z)|^2}{S(z, z)}$$

where $D = \{(z_1, z_2) | \mathrm{Im} z_1 - \Phi(z_2, z_2) \in \Omega\}$ and B is the distinguished boundary of D:

$$B = \{(z_1, z_2) | \mathrm{Im} z_1 - \Phi(z_2, z_2) = 0\}.$$

The Szegö kernel function S is given by

$$S(z, w) = \frac{1}{(2\pi)^{n_1}} \int_{\Omega'} e^{-<\alpha, \rho(z,w)>} L(\alpha)^{-1} d\alpha,$$

where

$$\rho(z, w) = \frac{z_1 - \overline{w_1}}{i} - 2\Phi(z_2, w_2)$$

for $z = (z_1, z_2)$, $w = (w_1, w_2)$, and

$$L(\alpha) = (\frac{i}{2})^{n_2} \int_{V_2} e^{-2<\alpha, \Phi(z_2, z_2)>} dz_2 d\overline{z_2}.$$

Now, we define the Poisson kernel as follows:

$$P(Z, W) = \frac{\rho(Z)}{|\rho(Z, W)|^2}, \quad \text{for} \quad Z = (z_1, z_2) \in \mathfrak{S}_2, W = (w_1, w_2) \in \partial \mathfrak{S}_2, \tag{3.1}$$

where

$$\rho(Z, W) := \overline{z_1} + w_1 - \overline{z_2} w_2.$$

In fact, for $Z = (\frac{\rho + |z|^2}{2} + it, z)$ and $Z' = (\frac{\rho' + |z'|^2}{2} + it', z')$, where $\rho > 0$, $\rho' > 0$, $t, t' \in \mathbb{R}$ and $z, z' \in \mathbb{C}$, we have

$$|\rho(Z, Z')|^2 = \frac{1}{4}(\rho + \rho' + |z - z'|^2)^2 + (t' - t + \mathrm{Im} z \overline{z'})^2 \geq 0.$$

$\rho(Z, Z') = 0$ if and only if $Z = Z' \in \partial\mathfrak{S}_2$.

In their papers [Ko], [KW], Korányi and Wolf studied the realization of Siegel domain of type II and the Poisson integral. Our definition has a little difference from theirs. We have

$$LP(Z,W)^s = s(s-2)P(Z,W)^s - s^2\rho(W)P(Z,W)^{s+1} = s(s-2)P(Z,W)^s. \tag{3.2}$$

In fact, by Helgason's conjecture, which was proved by Kashiwara et al. in [K], that the eigenfunctions on Riemannian symmetric spaces can be represented as Poisson integrals of their hyperfunction boundary values.

The boundary of $\mathfrak{S}_2$:

$$\partial\mathfrak{S}_2 = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 + \overline{z_1} = z_2\overline{z_2}\}. \tag{3.3}$$

In the homogeneous coordinates, one has

$$\overline{\partial\mathfrak{S}_2} = \{(z_1, z_2, z_3) \in \mathbb{C}^3 : z_1\overline{z_3} + \overline{z_1}z_3 = z_2\overline{z_2}\}.$$

In particular,

$$\overline{\partial\mathfrak{S}_2}(\mathcal{O}) = \{(a_1, a_2, a_3) \in \mathcal{O}^3 : a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}\}.$$

The Green function associated to $Lf = 0$ on $\mathfrak{S}_2$ is defined as

$$G(Z,W) = \log|\rho(Z,W)| \quad \text{for} \quad Z, W \in \mathfrak{S}_2. \tag{3.4}$$

It satisfies that $LG(Z,W) = 0$.

If $g = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in U(2,1)$, then $g^*Jg = J$, i.e.,

$$\begin{cases} -\overline{c_1}a_1 + \overline{b_1}b_1 - \overline{a_1}c_1 = 0, \\ -\overline{c_2}a_1 + \overline{b_2}b_1 - \overline{a_2}c_1 = 0, \\ -\overline{c_2}a_2 + \overline{b_2}b_2 - \overline{a_2}c_2 = 1, \\ -\overline{c_3}a_1 + \overline{b_3}b_1 - \overline{a_3}c_1 = -1, \\ -\overline{c_3}a_2 + \overline{b_3}b_2 - \overline{a_3}c_2 = 0, \\ -\overline{c_3}a_3 + \overline{b_3}b_3 - \overline{a_3}c_3 = 0. \end{cases}$$

By a straightforward calculation, we have

$$\rho(g(Z), g(W)) = \frac{\overline{z_1} + w_1 - \overline{z_2}w_2}{\overline{(c_1z_1 + c_2z_2 + c_3)}(c_1w_1 + c_2w_2 + c_3)}.$$

Hence,

$$\rho(g(Z), g(W))\overline{j(g, Z)}j(g, W) = \rho(Z, W), \quad \text{for} \quad g \in G. \tag{3.5}$$

Therefore,

$$P(g(Z), g(W)) = |j(g, W)|^2 P(Z, W). \tag{3.6}$$

By Proposition 2.9, we have

$$P(g(Z), W) = |j(g, g^{-1}(W))|^2 P(Z, g^{-1}(W)) = |j(g^{-1}, W)|^{-2} P(Z, g^{-1}(W)). \tag{3.7}$$

The point-pair invariant on $\mathfrak{S}_2$ is defined by

$$\sigma(Z, Z') := \frac{|\rho(Z, Z')|^2}{\rho(Z)\rho(Z')}, \tag{3.8}$$

which is invariant under the action of $U(2,1)$. The critical exponent is given by

$$\delta(\Gamma) = \inf\{s > 0 : \sum_{\gamma\in\Gamma} \sigma(Z, \gamma(Z'))^{-s} < \infty\}. \tag{3.9}$$

The number $\delta(\Gamma)$ does not depend on the choice of Z, Z'. It is an invariant of the group Γ.

The Eisenstein series is given by

$$E(Z, W; s) := \sum_{\gamma\in\Gamma} P(\gamma(Z), W)^s, \quad \text{for} \quad \mathrm{Re}(s) > \delta(\Gamma), \tag{3.10}$$

where Γ is a discrete subgroup of $U(2,1)$. We have the following lemma:

LEMMA 3.1. *The properties of Eisenstein series:*

1. $LE(Z, W; s) = s(s-2)E(Z, W; s)$.
2. $E(\gamma(Z), W; s) = E(Z, W; s)$.
3. $E(Z, \gamma(W); s) = |j(\gamma, W)|^{2s} E(Z, W; s)$ *for* $\gamma \in \Gamma$.

We have the following lemma:

LEMMA 3.2. *Let* $f = f(u)$ *with* $u = u(Z, Z') = \sigma(Z, Z') - 1$, *then*

$$Lf = u(u+1)f''(u) + (3u+2)f'(u). \tag{3.11}$$

By $Lf = s(s-2)f$, we have

$$u(u+1)f''(u) + (3u+2)f'(u) + s(2-s)f(u) = 0. \tag{3.12}$$

A solution is

$$\varphi_s(u(Z,Z')) = \frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s-\frac{1}{2})}(\frac{1}{4})^{s-1}u^{-s}F(s,s-1;2s-1;-u^{-1}). \tag{3.13}$$

Here $F = {}_2F_1$ is the Gauss hypergeometric function.

Let

$$G(Z,Z';s) = \sum_{\gamma\in\Gamma} r(Z,\gamma(Z');s), \quad \text{for} \quad \mathrm{Re}(s) > \delta(\Gamma), \tag{3.14}$$

where $r(Z,Z';s) = \varphi_s(u(Z,Z'))$. By the integral representation of hypergeometric functions, we can only consider the series

$$\sum_{\gamma\in\Gamma} \frac{1}{[1+u(Z,\gamma(Z'))]^s} = \sum_{\gamma\in\Gamma} \sigma(Z,\gamma(Z'))^{-s},$$

which is convergent for $\mathrm{Re}(s) > \delta(\Gamma)$. Thus, the right hand side of (3.14) is convergent when $\mathrm{Re}(s) > \delta(\Gamma)$. We have

$$\lim_{\rho(Z)\to 0} \rho(Z)^{-s} r(Z,Z';s) = \frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s-\frac{1}{2})}(\frac{1}{4})^{s-1}P(Z',W)^s,$$

where $Z \to W \in \partial\mathfrak{S}_2$ as $\rho(Z) \to 0$. Consequently,

$$\lim_{\rho(Z)\to 0} \rho(Z)^{-s} G(Z,Z';s) = \frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s-\frac{1}{2})}(\frac{1}{4})^{s-1}E(Z',W;s).$$

In the real hyperbolic geometry, the double transitivity of the action of the isometric groups on the real hyperbolic space has been studied extensively (see [ElGM] for three dimensional hyperbolic space). Now, we give the corresponding results on the complex hyperbolic space.

PROPOSITION 3.3. *The group $U(2,1)$ acts in the following sense doubly transitively on $\mathfrak{S}_2$: For all $P, P', Q, Q' \in \mathfrak{S}_2$, such that $d(P,P') = d(Q,Q')$, there exists an element $g \in U(2,1)$ such that $g(P) = Q$, $g(P') = Q'$.*

Proof. Set $P = (z_1, z_2)$ with $z_1 = -\omega\rho + \frac{|z|^2}{2} + it$, $z_2 = z$, where $\rho > 0$, $t \in \mathbb{R}$ and $z \in \mathbb{C}$. Then $\rho(z_1, z_2) = \rho$.

$$T_1 := \begin{pmatrix} \frac{1}{\sqrt{\rho}} & & \\ & 1 & \\ & & \sqrt{\rho} \end{pmatrix} \begin{pmatrix} 1 & -\overline{z} & \frac{|z|^2}{2} - it \\ & 1 & -z \\ & & 1 \end{pmatrix} \in U(2,1).$$

Moreover, $T_1(z_1, z_2) = (-\omega, 0)$, i.e. T_1 maps P onto $(-\omega, 0)$. Set $(z_1', z_2') = (-\omega\rho' + \frac{|z'|^2}{2} + it', z')$, then

$$T_1(P') = \left(-\omega\frac{\rho'}{\rho} + \frac{1}{2}\left(\frac{|z'-z|}{\sqrt{\rho}}\right)^2 + i\frac{t'-t+\operatorname{Im}(z\overline{z'})}{\rho}, \frac{z'-z}{\sqrt{\rho}}\right),$$

i.e., $T_1(P')$ has the form of $(-\omega\rho + \frac{|z|^2}{2} + it, z)$.

If $k(-\omega\rho + \frac{|z|^2}{2} + it, z) = (-\omega\rho', 0)$, where $k \in K$, then we obtain two complex equations about $\alpha, \beta, \gamma, \delta$ and θ. In fact, the number of real linear equations is four. While, the real dimension of K is five. Hence, the existence of k is proved. Applying a suitable element T_2 of the stabilizer K of $(-\omega, 0)$ in $U(2,1)$ to $T_1(P')$ we get a transform $T = T_2T_1$ such that $T(P) = (-\omega, 0)$, $T(P') = (-\lambda\omega, 0)$ with $\lambda \geq 1$. Since the isometric group of $\mathfrak{S}_2$ is $U(2,1)$, we have

$$d(P, P') = d(T(P), T(P')) = d((-\omega, 0), (-\lambda\omega, 0)) = \int_1^\lambda \frac{dr}{r} = \log\lambda.$$

So $\lambda = \exp(d(P, P'))$. Similarly, there exists an element $S \in U(2,1)$ such that $S(Q) = (-\omega, 0)$, $S(Q') = (-\lambda\omega, 0)$ with the same λ, since $d(P, P') = d(Q, Q')$. Thus, $g := S^{-1}T$ has the desired properties.

$\square$

In [ElGM], the concept of real hyperbolic distance was introduced and the basic properties was studied. Now, we give the complex hyperbolic distance and obtain the triangle inequality for our function δ.

Proposition 3.4. *The complex hyperbolic distance $d(P, P')$ is given by*

$$\cosh d(P, P') = \delta(P, P'), \tag{3.15}$$

where δ is defined by

$$\delta(P, P') = \delta(Z, Z') = \frac{1}{2}[\sigma(Z, Z') + 1], \tag{3.16}$$

and $P = Z$, $P' = Z'$.

Proof. If $P = (-\omega, 0)$ and $P' = (-\lambda\omega, 0)$ $(\lambda \geq 1)$, we have

$$d((-\omega, 0), (-\lambda\omega, 0)) = \int_1^\lambda \frac{dr}{r} = \log\lambda.$$

$$\delta((-\omega, 0), (-\lambda\omega, 0)) = \frac{1}{2}(\lambda + \frac{1}{\lambda}) = \cosh d((-\omega, 0), (-\lambda\omega, 0)).$$

Thus the proposition is true in the special case $P=(-\omega,0)$, $P'=(-\lambda\omega,0)$. Note that δ is a point-pair invariant. Since $U(2,1)$ acts doubly transitively on $\mathfrak{S}_2$, there exists for all $P,P'\in\mathfrak{S}_2$ an element $g\in U(2,1)$, such that $g(P)=(-\omega,0)$, $g(P')=(-\lambda\omega,0)$, $\lambda=\exp(d(P,P'))$. Therefore, by point-pair invariance, we have

$$\begin{aligned}\cosh d(P,P')=&\cosh d((-\omega,0),(-\lambda\omega,0))=\delta((-\omega,0),(-\lambda\omega,0))=\delta(g(P),g(P'))\\=&\delta(P,P').\end{aligned}$$

□

Proposition 3.5. *For* $g=\begin{pmatrix}a_1&a_2&a_3\\b_1&b_2&b_3\\c_1&c_2&c_3\end{pmatrix}\in U(2,1)$*, the following formulas hold:*

(1)

$$\begin{aligned}&\delta(P,g(P))+\delta(Q,g(Q))+\delta(P,g(Q))+\delta(Q,g(P))\\=&\frac{5}{2}+2(|a_1|^2+|a_3|^2+|c_1|^2+|c_3|^2)+(|a_2|^2+|c_2|^2+|b_1|^2+|b_3|^2)+\frac{1}{2}|b_2|^2,\end{aligned}\tag{3.17}$$

where $P=(-\omega,0)$ *and* $Q=(-\overline{\omega},0)$*.*

(2)

$$\delta((1,0),g(1,0))=\frac{5}{8}+\frac{1}{8}\sum_{j=1}^{3}(|a_j|^2+|b_j|^2+|c_j|^2).\tag{3.18}$$

Proof. It is obtained by a straightforward calculation.

□

Proposition 3.6. *The triangle inequality for the metric function* δ *on the complex hyperbolic space: For all* $P,Q,R\in\mathfrak{S}_2$*, the following inequality holds:*

$$\frac{1}{72}\frac{\delta(Q,R)}{\delta(P,Q)}\leq\delta(P,R)\leq 72\delta(P,Q)\delta(Q,R).\tag{3.19}$$

Proof. Since δ is a point-pair invariant, it is sufficient to prove (3.19) only in the special case $Q=(\frac{1}{2},0)$. Set $P=(\frac{\rho_1+|z_1|^2}{2}+it_1,z_1)$ and $R=(\frac{\rho_2+|z_2|^2}{2}+it_2,z_2)$. Then

$$\delta(P,Q)=\frac{\frac{1}{4}(1+\rho_1+|z_1|^2)^2+t_1^2+\rho_1}{2\rho_1},\quad\delta(Q,R)=\frac{\frac{1}{4}(1+\rho_2+|z_2|^2)^2+t_2^2+\rho_2}{2\rho_2},$$

and

$$\delta(P,R)=\frac{1}{8\rho_1\rho_2}[(\rho_1+\rho_2+|z_1-z_2|^2)^2+4(t_2-t_1+\mathrm{Im}z_1\overline{z_2})^2+4\rho_1\rho_2].$$

Note that $|z_1 - z_2|^2 \leq 2(|z_1|^2 + |z_2|^2)$ and $|\mathrm{Im} z_1 \overline{z_2}| \leq |z_1||z_2| \leq \frac{1}{2}(|z_1|^2 + |z_2|^2)$, we have

$$8\rho_1\rho_2\delta(P,R) \leq (\rho_1 + \rho_2 + 2|z_1|^2 + 2|z_2|^2)^2 + (2|t_1| + 2|t_2| + |z_1|^2 + |z_2|^2)^2 + 4\rho_1\rho_2.$$

By $2|t_1||t_2| \leq t_1^2 + t_2^2$ and $2|t_i||z_j|^2 \leq t_i^2 + |z_j|^4$, $i, j = 1, 2$, we have

$$\begin{aligned}
&8\rho_1\rho_2\delta(P,R)\\
\leq&\rho_1^2 + \rho_2^2 + 9|z_1|^4 + 9|z_2|^4 + 6\rho_1\rho_2 + 4(\rho_1 + \rho_2)(|z_1|^2 + |z_2|^2) + 10|z_1|^2|z_2|^2\\
&+ 12t_1^2 + 12t_2^2\\
\leq&9(1 + \rho_1^2 + |z_1|^4 + 6\rho_1 + 2|z_1|^2 + 2\rho_1|z_1|^2 + 4t_1^2)\\
&\times (1 + \rho_2^2 + |z_2|^4 + 6\rho_2 + 2|z_2|^2 + 2\rho_2|z_2|^2 + 4t_2^2)\\
=&576\delta(P,Q)\delta(Q,R).
\end{aligned}$$

Hence, $\delta(P,R) \leq 72\delta(P,Q)\delta(Q,R)$. Interchanging P and Q, we obtain the other part of the inequality.

□

THEOREM 3.7. *Let*

$$\phi(Z, Z'; s) = \int_{\partial\mathfrak{S}_2} P(Z,W)^s P(Z',W)^{2-s} dm(W). \tag{3.20}$$

Then one has the following formula:

$$\phi(Z,Z';s) = -\frac{\pi}{s-1}\frac{\sqrt{\pi}\Gamma(\frac{3}{2}-s)}{\Gamma(2-s)}(\frac{1}{4})^{s-1}u^{-s}F(s, s-1; 2s-1; -u^{-1}) + (s \mapsto 2-s), \tag{3.21}$$

where $dm(W)$ is the Lebesque measure on $\partial\mathfrak{S}_2$, $u = u(Z,Z')$ is the point-pair invariant, and $(s \mapsto 2-s)$ is equal to the first term on the right hand side with s replaced by $2-s$.

Proof. $\phi(Z,Z';s)$ is a point-pair invariant. Thus, it suffices to calculate it for $Z = (z,t,\rho) = (0,0,\rho)$ and $Z' = (z',t',\rho') = (0,0,\rho')$, i.e., $Z = (\frac{1}{2}\rho, 0)$ and $Z' = (\frac{1}{2}\rho', 0)$. Let ρ and ρ' stand for $(\rho,0)$ and $(\rho',0)$, respectively. Now, $\rho(Z,Z') = \frac{1}{2}(\rho+\rho')$ and

$$\sigma(Z,Z') = \frac{(\rho+\rho')^2}{4\rho\rho'} = \frac{(\lambda+1)^2}{4\lambda}, \quad u = \frac{(\lambda-1)^2}{4\lambda},$$

where $\lambda = \rho/\rho'$. For $W \in \partial\mathfrak{S}_2$, $w_1 = \frac{1}{2}|w|^2 + iv$ and $w_2 = w$, where $w \in \mathbb{C}$, $v \in \mathbb{R}$. $\rho(Z,W) = \frac{1}{2}(\rho + |w|^2) + iv$ and $\rho(Z',W) = \frac{1}{2}(\rho' + |w|^2) + iv$. Hence,

$$\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; s) = \int_{\mathbb{C}}\int_{\mathbb{R}} \frac{\rho^s {\rho'}^{2-s}}{|\frac{1}{2}(\rho + |w|^2) + iv|^{2s}|\frac{1}{2}(\rho' + |w|^2) + iv|^{2(2-s)}} dm(w)dv.$$

By transform $w = re^{i\theta}$ with $r \geq 0, \theta \in [0, 2\pi)$ and $u = r^2$, we have

$$\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; s) = 16\pi\rho^s {\rho'}^{2-s} \int_0^\infty \int_0^\infty \frac{dudv}{[(\rho+u)^2+v^2]^s[(\rho'+u)^2+v^2]^{2-s}}.$$

Set $v = (\rho'+u)\sqrt{w}$, then

$$\int_0^\infty \frac{dv}{[(\rho+u)^2+v^2]^s[(\rho'+u)^2+v^2]^{2-s}} = \frac{1}{2}(\rho'+u)^{-3}\int_0^\infty w^{-\frac{1}{2}}(1+w)^{s-2}(l^2+w)^{-s}dw,$$

where $l = l(u) = \frac{\rho+u}{\rho'+u}$.

By (see [Er], p.115, (5))

$$F(a,b;c;1-z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)}\int_0^\infty s^{b-1}(1+s)^{a-c}(1+sz)^{-a}ds$$

for $\mathrm{Re}(c) > \mathrm{Re}(b) > 0$, $|\arg(z)| < \pi$, we have

$$\int_0^\infty w^{-\frac{1}{2}}(1+w)^{s-2}(1+wl^{-2})^{-s}dw = \frac{\pi}{2}F(s, \frac{1}{2}; 2; 1-l^{-2}).$$

Now, we need the following two formulas:

(1) $z = 1-l^{-2}$, $\frac{z}{z-1} = 1-l^2$.

By (see [Er], p.105, 2.9.(3))

$$F(a,b;c;z) = (1-z)^{-a}F(a, c-b; c; \frac{z}{z-1}),$$

one has

$$F(s, \frac{1}{2}; 2; 1-l^{-2}) = l^{2s}F(s, \frac{3}{2}; 2; 1-l^2).$$

(2) $z = 1-l^2$, $1-z = l^2$.

By (see [Er], p.108, 2.10.(1))

$$\begin{aligned} F(a,b;c;z) = {} & \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}F(a,b;a+b-c+1;1-z) \\ & + \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)}(1-z)^{c-a-b}F(c-a, c-b; c-a-b+1; 1-z), \end{aligned}$$

where $|\arg(1-z)| < \pi$, one has

$$\begin{aligned} F(s, \frac{3}{2}; 2; 1-l^2) = {} & \frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}F(s, \frac{3}{2}; s+\frac{1}{2}; l^2) \\ & + \frac{2\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}l^{1-2s}F(2-s, \frac{1}{2}; \frac{3}{2}-s; l^2). \end{aligned}$$

Hence, $\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; s) = \phi_I + \phi_{II}$, where

$$\phi_I = 4\pi^2\lambda^s{\rho'}^2\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}\int_0^\infty(\rho'+u)^{-3}F(s,\frac{3}{2};s+\frac{1}{2};l^2)du,$$

$$\phi_{II} = 4\pi^2\lambda^s{\rho'}^2\frac{2\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}\int_0^\infty(\rho'+u)^{-3}l^{1-2s}F(2-s,\frac{1}{2};\frac{3}{2}-s;l^2)du.$$

$l = \frac{\rho+u}{\rho'+u}$ implies that $u = \frac{\rho-\rho' l}{l-1}$. Thus, $du = \frac{\rho'-\rho}{(l-1)^2}dl$ and $\rho'+u = \frac{\rho-\rho'}{l-1}$. Without loss of generality, we can assume that $\rho < \rho'$. Then

$$\phi_I = -\frac{4\pi^2\lambda^s}{(1-\lambda)^2}\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}\int_\lambda^1(l-1)F(s,\frac{3}{2};s+\frac{1}{2};l^2)dl,$$

$$\phi_{II} = -\frac{4\pi^2\lambda^s}{(1-\lambda)^2}\frac{2\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}\int_\lambda^1(l-1)l^{1-2s}F(2-s,\frac{1}{2};\frac{3}{2}-s;l^2)dl.$$

We have

$$\begin{aligned}&\int_\lambda^1(l-1)F(s,\frac{3}{2};s+\frac{1}{2};l^2)dl\\=&\sum_{k=0}^\infty\frac{(s)_k(\frac{3}{2})_k}{k!(s+\frac{1}{2})_k}(\frac{\lambda^{2k+1}}{2k+1}-\frac{\lambda^{2k+2}}{2k+2}-\frac{1}{(2k+1)(2k+2)})\\=&\lambda F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)-\frac{2s-1}{2s-2}F(s-1,\frac{1}{2};s-\frac{1}{2};\lambda^2)+\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}.\end{aligned}$$

Here we use the following formula:

$$F(s-1,-\frac{1}{2};s-\frac{1}{2};1) = \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}, \quad \text{for} \quad \mathrm{Re}(s) > 0.$$

Similarly,

$$\begin{aligned}&\int_\lambda^1(l-1)l^{1-2s}F(2-s,\frac{1}{2};\frac{3}{2}-s;l^2)dl\\=&\frac{\lambda^{2-2s}}{2-2s}F(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2)-\frac{\lambda^{3-2s}}{3-2s}F(2-s,\frac{1}{2};\frac{5}{2}-s;\lambda^2)-\frac{1}{2-2s}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)},\end{aligned}$$

for $\mathrm{Re}(s) < 2$.

Set $z=-\lambda$, then $\frac{4z}{(1+z)^2}=-u^{-1}$. By (see [Er], p.111, 2.11.(5))

$$F(a,b;2b;\frac{4z}{(1+z)^2})=(1+z)^{2a}F(a,a+\frac{1}{2}-b;b+\frac{1}{2};z^2),$$

we have

$$F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)=(1-\lambda)^{-2s}F(s,s;2s;-u^{-1}),$$

$$F(s-1,\frac{1}{2};s-\frac{1}{2};\lambda^2)=(1-\lambda)^{2-2s}F(s-1,s-1;2s-2;-u^{-1}),$$

$$F(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2)=(1-\lambda)^{2s-2}F(1-s,1-s;2-2s;-u^{-1}),$$

$$F(2-s,\frac{1}{2};\frac{5}{2}-s;\lambda^2)=(1-\lambda)^{2s-4}F(2-s,2-s;4-2s;-u^{-1}).$$

Thus,

$$\begin{aligned}\phi_I=&-4\pi^2\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}(4u)^{-s-1}F(s,s;2s;-u^{-1})\\&-\frac{4\pi^2}{s-1}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}(4u)^{-s}F(s-1,s-1;2s-2;-u^{-1})\\&-\frac{4\pi}{s-1}\frac{\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2})}{\Gamma(s)\Gamma(2-s)}\lambda^s(1-\lambda)^{-2},\end{aligned}$$

and

$$\begin{aligned}\phi_{II}=&-\frac{4\pi^2}{1-s}\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}(4u)^{s-2}F(1-s,1-s;2-2s;-u^{-1})\\&-4\pi^2\frac{\Gamma(s-\frac{3}{2})}{\sqrt{\pi}\Gamma(s)}(4u)^{s-3}F(2-s,2-s;4-2s;-u^{-1})\\&-\frac{4\pi}{s-1}\frac{\Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)}{\Gamma(s)\Gamma(2-s)}\lambda^s(1-\lambda)^{-2}.\end{aligned}$$

Note that

$$\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2})+\Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)=(s-\frac{1}{2})\Gamma(\frac{1}{2}-s)\Gamma(s-\frac{1}{2})+(\frac{1}{2}-s)\Gamma(s-\frac{1}{2})\Gamma(\frac{1}{2}-s)=0.$$

Moreover, we need the following lemma.

Lemma 3.8. *The following formula about hypergeometric functions holds:*

$$\gamma F-\beta zF(\beta+1,\gamma+1)-\gamma F(\alpha-1)=0,$$

where F denotes $F(\alpha,\beta;\gamma;z)$, $F(\beta+1,\gamma+1)$ stands for $F(\alpha,\beta+1;\gamma+1;z)$ and $F(\alpha-1)$ stands for $F(\alpha-1,\beta;\gamma;z)$.

Proof. It is obtained by the integral representation of hypergeometric functions.

□

By Lemma 3.8 and the following formula (see [Er], p.103, (35)):

$$(\gamma-1)F(\gamma-1)-\alpha F(\alpha+1)-(\gamma-\alpha-1)F=0.$$

we get

$$(2s-1)F(s,s-1;2s-1;z)-(s-1)zF(s,s;2s;z)-(2s-1)F(s-1,s-1;2s-1;z)=0,$$

$$(2s-2)F(s-1,s-1;2s-2;z)-(s-1)F(s,s-1;2s-1;z)-(s-1)F(s-1,s-1;2s-1;z)=0.$$

Thus, we have

$$\frac{4s-2}{s-1}F(s,s-1;2s-1;z)=zF(s,s;2s;z)+\frac{4s-2}{s-1}F(s-1,s-1;2s-2;z).$$

Substituting this identity to the expression of $\phi(\frac{1}{2}\rho,\frac{1}{2}\rho';s)$, We get the desired results.

□

Consequently, we have

Corollary 3.9. *The following formula holds:*

$$\phi(Z,Z';s)=-\frac{\pi}{s-1}c(s)c(2-s)[r(Z,Z';s)-r(Z,Z';2-s)], \tag{3.22}$$

where

$$c(s)=\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}.$$

Corollary 3.10. *The functional equation for the trivial group on $\mathfrak{S}_2$: If Re$(s)>1$, then the following identity holds:*

$$\int_{\partial\mathfrak{S}_2}P(Z,W)^s|\rho(W,W')|^{-2(2-s)}dm(W)=\frac{\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}4^{s-1}P(Z,W')^{2-s}, \tag{3.23}$$

where $W'\in\partial\mathfrak{S}_2$.

Proof. The integral on the left-hand side is absolutely convergent. We set

$$P(Z',W)=\frac{\rho(Z')}{|\rho(Z',W)|^2}$$

in the formula of Theorem 3.7. Next, we multiply both sides of the formula by $\rho(Z')^{s-2}$ and take the limit as $Z'\to W'\in\partial\mathfrak{S}_2$. Thus we get the desired results.

□

Proposition 3.11. *The integral formula for the Poisson kernel:*

$$\int_{\partial\mathfrak{S}_2} P(Z,W)^s dm(W) = \frac{\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}4^{s-1}\rho^{2-s}, \quad \textit{for} \quad Re(s)>1. \tag{3.24}$$

Proof. Denote the left hand side of (3.24) as $f(Z)$, then

$$f(Z) = \rho^s \int_{\partial\mathfrak{S}_2} |\overline{z_1} + w_1 - \overline{z_2}w_2|^{-2s} dm(W).$$

Set $z_1 = \frac{\rho+|z|^2}{2} + ir$, $z_2 = z$, $w_1 = \frac{|w|^2}{2} + it$ and $w_2 = w$, where $z, w \in \mathbb{C}$, $t, r \in \mathbb{R}$, we have

$$\begin{aligned} f(Z) &= \rho^s \int_{\mathbb{C}}\int_{\mathbb{R}} |\frac{\rho+|w-z|^2}{2} + i(t-r+\mathrm{Im}(z\overline{w}))|^{-2s} dm(w)dt \\ &= \rho^s \int_{\mathbb{C}}\int_{\mathbb{R}} |\frac{\rho+|w|^2}{2} + it|^{-2s} dm(w)dt \\ &= 2\pi\rho^s \int_0^\infty \int_{-\infty}^\infty \frac{rdrdt}{[\frac{1}{4}(\rho+r^2)^2+t^2]^s} \\ &= 2^{2s-1}\pi\rho^s \int_0^\infty \int_{-\infty}^\infty \frac{drdu}{[(\rho+r)^2+u^2]^s}. \end{aligned}$$

By the well known formula (see [Ku], p.15):

$$\int_{-\infty}^\infty \frac{dt}{(1+t^2)^s} = \sqrt{\pi}\frac{\Gamma(s-\frac{1}{2})}{\Gamma(s)}, \quad \text{for} \quad \mathrm{Re}(s)>1,$$

we have

$$f(Z) = \frac{\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}4^{s-1}\rho^{2-s}.$$

□

We define the S-matrix as follows:

$$S(W,W';s) := \sum_{\gamma\in\Gamma} |\rho(\gamma(W),W')|^{-2s}|j(\gamma,W)|^{-2s} \tag{3.25}$$

for $W, W' \in \Omega(\Gamma)$ and $\mathrm{Re}(s) > \delta(\Gamma)$.

Theorem 3.12. *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following functional equation for Eisenstein series holds:*

$$\int_{\Gamma\backslash\Omega(\Gamma)} E(Z,W;s)S(W,W';2-s)dm(W) = \frac{4^{s-1}\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}E(Z,W';2-s), \tag{3.26}$$

for $1 < Re(s) < 2 - \delta(\Gamma)$.

Proof. By Corollary 3.10,

$$\int_{\partial\mathfrak{S}_2} E(Z,W;s)|\rho(W,W')|^{-2(2-s)}dm(W) = \frac{4^{s-1}\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}E(Z,W';2-s). \tag{3.27}$$

On the other hand,

$$\begin{aligned}\int_{\partial\mathfrak{S}_2} f(W)dW &= \int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma} f(\gamma(W))dm(\gamma(W))\\ &= \int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma}|J(\gamma(W))|f(\gamma(W))dm(W).\end{aligned}$$

Hence, the left hand side of (3.27):

$$\begin{aligned}LHS &= \int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma}|J(\gamma(W))|E(Z,\gamma(W);s)|\rho(\gamma(W),W')|^{-2(2-s)}dm(W)\\ &= \int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma}|j(\gamma,W)|^{-4}|j(\gamma,W)|^{2s}E(Z,W;s)|\rho(\gamma(W),W')|^{-2(2-s)}dm(W)\\ &= \int_{\Gamma\backslash\Omega(\Gamma)} E(Z,W;s)S(W,W';2-s)dm(W).\end{aligned}$$

□

PROPOSITION 3.13. *Assume that* Γ *is convex cocompact and* $\delta(\Gamma) < 1$. *Then the following product formula holds:*

$$\begin{aligned}&\int_{\Gamma\backslash\Omega(\Gamma)} E(Z,W;s)E(Z',W;2-s)dm(W)\\ &= -\frac{\pi}{s-1}c(s)c(2-s)[G(Z,Z';s)-G(Z,Z';2-s)],\end{aligned} \tag{3.28}$$

where $\delta(\Gamma) < Re(s) < 2 - \delta(\Gamma)$.

For $s \in \mathbb{C}$, similar as in [Ma1] and [Ma2], we set

$$L_s^2(\Gamma) := \{f(s,\cdot) : f(s,\cdot) \in C^\infty(\Omega(\Gamma)), f(s,\gamma(W))j(\gamma,W)^{2-s} = f(s,W), \gamma\in\Gamma\}.$$

If $f(s,\cdot) \in L_s^2(\Gamma)$, we define the Eisenstein integral:

$$E(Z,\cdot,s) : L_s^2(\Gamma) \to C^\infty(\Gamma\backslash\mathfrak{S}_2),$$

$$E(Z,f,s)=\int_{\Gamma\backslash\Omega(\Gamma)}E(Z,W;s)f(s,W)dm(W),\quad \text{for}\quad \mathrm{Re}(s)>\delta(\Gamma), \tag{3.29}$$

and the scattering operator:

$$S_s(f)(z,t):=\lim_{Z\to(z,t)}\rho(Z)^{-s}[E(Z,f,s)-\frac{4^{s-1}\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}f(s,(z,t))\rho(Z)^{2-s}], \tag{3.30}$$

where $\delta(\Gamma)<\mathrm{Re}(s)<2$.

Now, the functional equation for Eisenstein series can be written in terms of Eisenstein integrals as

$$E(Z,S_{2-s}(f),s)=\frac{4^{s-1}\pi}{s-1}\frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)}E(Z,f,2-s), \tag{3.31}$$

and in terms involving the scattering operator as:

$$S_sS_{2-s}=-\frac{\pi^3}{(s-1)^2}\frac{\Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)}{\Gamma(s)\Gamma(2-s)}I. \tag{3.32}$$

4. Product formulas of weight k on $U(2,1)$

In the case of $SL(2,\mathbb{R})$, many people, especially Maass [M], Selberg [Se], Roelcke [Ro], Fay [Fa], Elstrodt [El], Hejhal [H], and Shimura [Sh2] studied the eigenvalue problems which concern a homogeneous equation

$$(\Delta_k-\lambda)f(z)=0.$$

Here f is a function on the complex upper plane

$$\mathbb{H}=\{z\in\mathbb{C}:\mathrm{Im}(z)>0\},$$

and L_k is a differential operator given by

$$\Delta_k=y^2\left(\frac{\partial^2}{\partial x^2}+\frac{\partial^2}{\partial y^2}\right)-iky\frac{\partial}{\partial x},$$

where $z=x+iy$ and k is an integer. Let Γ be a congruence subgroup of $SL(2,\mathbb{Z})$ and assume that f is a Γ-automorphic form of weight k in the sense that it is invariant under

$$f\mapsto f(\gamma(z))(cz+d)^{-k}|cz+d|^k$$

for any $\gamma = \begin{pmatrix} * & * \\ c & d \end{pmatrix} \in \Gamma$. In fact, L_k commutes with the above map. One of the most remarkable facts about such λ and f is the existence of Selberg zeta function (see [Se]), whose set of zeros coincides essentially with the set $S(\Gamma, k)$ of $s \in \mathbb{C}$ such that $\lambda = s(1-s)$ occurs as an eigenvalue of L_k as in the above equation with a cusp form f (see [Sh2]).

Now, we study the corresponding problems in the case of $U(2,1)$.

PROPOSITION 4.1. *Let*

$$\begin{aligned} L_{(\alpha,\beta)} &= (z_1 + \overline{z_1} - z_2\overline{z_2}) \\ &\times \left[(z_1+\overline{z_1})\frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2}\frac{\partial^2}{\partial z_1 \partial \overline{z_2}} + \beta \frac{\partial}{\partial z_1} + \alpha \frac{\partial}{\partial \overline{z_1}} \right] \end{aligned} \tag{4.1}$$

and

$$E(Z, \overline{Z}; \alpha, \beta) = \sum_{\gamma \in \Gamma \cap N \backslash \Gamma} j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta}, \tag{4.2}$$

where $\Gamma = U(2,1;\mathcal{O})$ *is the Picard modular group. Then the following formula holds:*

$$L_{(\alpha,\beta)} E(Z, \overline{Z}; \alpha, \beta) = 0. \tag{4.3}$$

Proof. We have

$$\begin{aligned} &\left[(z_1+\overline{z_1})\frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2}\frac{\partial^2}{\partial z_1 \partial \overline{z_2}} \right] [j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta}] \\ =&\alpha\beta[a_1 j(\gamma, Z)^{-\alpha-1} \overline{j(\gamma, Z)}^{-\beta} + \overline{a_1} j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta-1}] \\ =&-\left(\beta \frac{\partial}{\partial z_1} + \alpha \frac{\partial}{\partial \overline{z_1}} \right) [j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta}]. \end{aligned}$$

Hence, $L_{(\alpha,\beta)} E(Z, \overline{Z}; \alpha, \beta) = 0.$

□

Set

$$E_{3k}(Z, \overline{Z}; \beta) = \rho(Z)^{\beta} E(Z, \overline{Z}; \alpha, \beta) = \sum_{\gamma \in \Gamma \cap N \backslash \Gamma} \frac{\rho(Z)^{\beta}}{j(\gamma, Z)^{3k} |j(\gamma, Z)|^{2\beta}} \tag{4.4}$$

with $3k = \alpha - \beta$.

In particular, when $\beta = -k$ and $\alpha = k$, we have

$$L_k = (z_1 + \overline{z_1} - z_2\overline{z_2}) \times \left[(z_1 + \overline{z_1})\frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2}\frac{\partial^2}{\partial z_1 \partial \overline{z_2}} - k(\frac{\partial}{\partial z_1} - \frac{\partial}{\partial \overline{z_1}})\right].$$

Proposition 4.2. *The operator L_k has the following automorphic property:*

$$L_k[f(\gamma(Z))j(\gamma, Z)^{-k}\overline{j(\gamma, Z)}^k] = j(\gamma, Z)^{-k}\overline{j(\gamma, Z)}^k(L_k f)(\gamma(Z)), \tag{4.5}$$

for $\gamma \in U(2,1)$ and $Z = (z_1, z_2) \in \mathbb{C}^2$, where $f = f(Z, \overline{Z})$ is a real analytic function. In other words, L_k commutes with the following map

$$f(Z) \mapsto f(\gamma(Z))j(\gamma, Z)^{-k}\overline{j(\gamma, Z)}^k. \tag{4.6}$$

Proof. For $\gamma = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in U(2,1)$, one has $\gamma J \gamma^* = J$, i.e.,

$$\begin{cases} -\overline{a_1}a_3 + a_2\overline{a_2} - a_1\overline{a_3} = 0, \\ -a_3\overline{b_1} + a_2\overline{b_2} - a_1\overline{b_3} = 0, \\ -a_3\overline{c_1} + a_2\overline{c_2} - a_1\overline{c_3} = -1, \\ -b_3\overline{b_1} + b_2\overline{b_2} - b_1\overline{b_3} = 1, \\ -b_3\overline{c_1} + b_2\overline{c_2} - b_1\overline{c_3} = 0, \\ -c_3\overline{c_1} + c_2\overline{c_2} - c_1\overline{c_3} = 0. \end{cases}$$

Denote

$$\gamma(z_1, z_2) = (u(z_1, z_2), v(z_1, z_2)) = \left(\frac{a_1z_1 + a_2z_2 + a_3}{c_1z_1 + c_2z_2 + c_3}, \frac{b_1z_1 + b_2z_2 + b_3}{c_1z_1 + c_2z_2 + c_3}\right).$$

Set

$$A_{1,1} := \frac{\partial}{\partial z_1}\left(\frac{a_1z_1 + a_2z_2 + a_3}{c_1z_1 + c_2z_2 + c_3}\right) = [(a_1c_2 - a_2c_1)z_2 + (a_1c_3 - a_3c_1)]j(\gamma, Z)^{-2}.$$

$$A_{1,2} := \frac{\partial}{\partial z_2}\left(\frac{a_1z_1 + a_2z_2 + a_3}{c_1z_1 + c_2z_2 + c_3}\right) = [(a_2c_1 - a_1c_2)z_1 + (a_2c_3 - a_3c_2)]j(\gamma, Z)^{-2}.$$

$$A_{2,1} := \frac{\partial}{\partial z_1}\left(\frac{b_1z_1 + b_2z_2 + b_3}{c_1z_1 + c_2z_2 + c_3}\right) = [(b_1c_2 - b_2c_1)z_2 + (b_1c_3 - b_3c_1)]j(\gamma, Z)^{-2}.$$

$$A_{2,2} := \frac{\partial}{\partial z_2}\left(\frac{b_1 z_1 + b_2 z_2 + b_3}{c_1 z_1 + c_2 z_2 + c_3}\right) = [(b_2 c_1 - b_1 c_2) z_1 + (b_2 c_3 - b_3 c_2)] j(\gamma, Z)^{-2}.$$

We have

$$\begin{aligned} L_k(\gamma(Z)) =& (u + \overline{u} - v\overline{v}) \\ & \times \left[(u+\overline{u})\frac{\partial^2}{\partial u \partial \overline{u}} + \frac{\partial^2}{\partial v \partial \overline{v}} + v\frac{\partial^2}{\partial \overline{u} \partial v} + \overline{v}\frac{\partial^2}{\partial u \partial \overline{v}} - k\left(\frac{\partial}{\partial u} - \frac{\partial}{\partial \overline{u}}\right)\right], \end{aligned}$$

while $u + \overline{u} - v\overline{v} = (z_1 + \overline{z_1} - z_2\overline{z_2})|j(\gamma, Z)|^{-2}$.

On the other hand,

$$\begin{aligned} & L_k[f(\gamma(Z)) j(\gamma, Z)^{-k}\overline{j(\gamma, Z)}^k] \\ =& j(\gamma, Z)^{-k}\overline{j(\gamma, Z)}^k (z_1 + \overline{z_1} - z_2\overline{z_2})[F_{1,1}\frac{\partial^2 f}{\partial u \partial \overline{u}} + F_{1,2}\frac{\partial^2 f}{\partial u \partial \overline{v}} + F_{2,1}\frac{\partial^2 f}{\partial \overline{u} \partial v} \\ & + F_{2,2}\frac{\partial^2 f}{\partial v \partial \overline{v}} + H_1\frac{\partial f}{\partial u} + H_2\frac{\partial f}{\partial v} + G_1\frac{\partial f}{\partial \overline{u}} + G_2\frac{\partial f}{\partial \overline{v}}](\gamma(Z)), \end{aligned}$$

where

$$\begin{cases} F_{1,1} = (z_1 + \overline{z_1})A_{1,1}\overline{A_{1,1}} + A_{1,2}\overline{A_{1,2}} + z_2 A_{1,2}\overline{A_{1,1}} + \overline{z_2}A_{1,1}\overline{A_{1,2}}, \\ F_{1,2} = (z_1 + \overline{z_1})A_{1,1}\overline{A_{2,1}} + A_{1,2}\overline{A_{2,2}} + z_2 A_{1,2}\overline{A_{2,1}} + \overline{z_2}A_{1,1}\overline{A_{2,2}}, \\ F_{2,2} = (z_1 + \overline{z_1})A_{2,1}\overline{A_{2,1}} + A_{2,2}\overline{A_{2,2}} + z_2 A_{2,2}\overline{A_{2,1}} + \overline{z_2}A_{2,1}\overline{A_{2,2}}, \\ H_1 = -kA_{1,1} + k[(z_1 + \overline{z_1})\overline{c_1}A_{1,1} + \overline{c_2}A_{1,2} + z_2\overline{c_1}A_{1,2} + \overline{z_2 c_2}A_{1,1}] j(\overline{\gamma}, \overline{Z})^{-1}, \\ H_2 = -kA_{2,1} + k[(z_1 + \overline{z_1})\overline{c_1}A_{2,1} + \overline{c_2}A_{2,2} + z_2\overline{c_1}A_{2,2} + \overline{z_2 c_2}A_{2,1}] j(\overline{\gamma}, \overline{Z})^{-1}, \\ F_{2,1} = \overline{F_{1,2}}, \quad G_1 = -\overline{H_1}, \quad G_2 = -\overline{H_2}. \end{cases}$$

Now, we will prove that $F_{1,1} = (u + \overline{u})|j(\gamma, Z)|^{-2}$. Note that

$$\begin{aligned} & F_{1,1}|j(\gamma, Z)|^4 \\ & = (z_1 + \overline{z_1})[(a_1 c_2 - a_2 c_1) z_2 + (a_1 c_3 - a_3 c_1)][(\overline{a_1 c_2} - \overline{a_2 c_1})\overline{z_2} + (\overline{a_1 c_3} - \overline{a_3 c_1})] \\ & + [(a_2 c_1 - a_1 c_2) z_1 + (a_2 c_3 - a_3 c_2)][(\overline{a_2 c_1} - \overline{a_1 c_2})\overline{z_1} + (\overline{a_2 c_3} - \overline{a_3 c_2})] \\ & + z_2[(a_2 c_1 - a_1 c_2) z_1 + (a_2 c_3 - a_3 c_2)][(\overline{a_1 c_2} - \overline{a_2 c_1})\overline{z_2} + (\overline{a_1 c_3} - \overline{a_3 c_1})] \\ & + \overline{z_2}[(a_1 c_2 - a_2 c_1) z_2 + (a_1 c_3 - a_3 c_1)][(\overline{a_2 c_1} - \overline{a_1 c_2})\overline{z_1} + (\overline{a_2 c_3} - \overline{a_3 c_2})]. \end{aligned}$$

On the other hand,

$$u + \overline{u} = [(a_1 z_1 + a_2 z_2 + a_3)(\overline{c_1 z_1} + \overline{c_2 z_2} + \overline{c_3}) + (\overline{a_1 z_1} + \overline{a_2 z_2} + \overline{a_3})(c_1 z_1 + c_2 z_2 + c_3)]|j(\gamma, Z)|^{-2}.$$

Now, we need the following six identities:

$$\begin{aligned}
&(a_2c_1 - a_1c_2)(\overline{a_2c_1} - \overline{a_1c_2}) \\
=&(\overline{a_1}a_3 + a_1\overline{a_3})c_1\overline{c_1} - a_1\overline{a_2c_1}c_2 - \overline{a_1}a_2c_1\overline{c_2} + a_1\overline{a_1}(c_1\overline{c_3} + c_3\overline{c_1}) \\
=&\overline{a_1}c_1(a_3\overline{c_1} - a_2\overline{c_2} + a_1\overline{c_3}) + a_1\overline{c_1}(\overline{a_3}c_1 - \overline{a_2}c_2 + \overline{a_1}c_3) \\
=&\overline{a_1}c_1 + a_1\overline{c_1}.
\end{aligned}$$

$$\begin{aligned}
&(a_1c_2 - a_2c_1)(\overline{a_1c_3} - \overline{a_3c_1}) \\
=&\overline{a_1}c_2(1 - a_3\overline{c_1} + a_2\overline{c_2}) + a_2\overline{c_1}(1 + \overline{a_2}c_2 - \overline{a_1}c_3) - \overline{a_1}a_2c_1\overline{c_3} - a_1\overline{a_3c_1}c_2 \\
=&\overline{a_1}c_2 + a_2\overline{c_1} + \overline{c_1}c_2(-\overline{a_1}a_3 + a_2\overline{a_2} - a_1\overline{a_3}) + \overline{a_1}a_2(c_2\overline{c_2} - \overline{c_1}c_3 - c_1\overline{c_3}) \\
=&\overline{a_1}c_2 + a_2\overline{c_1}.
\end{aligned}$$

$$\begin{aligned}
&(a_2c_3 - a_3c_2)(\overline{a_1c_2} - \overline{a_2c_1}) + (a_1c_2 - a_2c_1)(\overline{a_2c_3} - \overline{a_3c_2}) \\
=&a_2\overline{c_2}(\overline{a_1}c_3 + \overline{a_3}c_1) + \overline{a_2}c_2(a_3\overline{c_1} + a_1\overline{c_3}) - \overline{a_1}a_3c_2\overline{c_2} - a_2\overline{a_2c_1}c_3 - a_2\overline{a_2}c_1\overline{c_3} - a_1\overline{a_3}c_2\overline{c_2} \\
=&a_2\overline{c_2} + \overline{a_2}c_2 + c_2\overline{c_2}(a_2\overline{a_2} - \overline{a_1}a_3 - a_1\overline{a_3}) + a_2\overline{a_2}(c_2\overline{c_2} - \overline{c_1}c_3 - c_1\overline{c_3}) \\
=&a_2\overline{c_2} + \overline{a_2}c_2.
\end{aligned}$$

$$\begin{aligned}
&(a_1c_3 - a_3c_1)(\overline{a_1c_3} - \overline{a_3c_1}) + (a_2c_1 - a_1c_2)(\overline{a_2c_3} - \overline{a_3c_2}) \\
=&a_1\overline{c_3}(\overline{a_1}c_3 - \overline{a_2}c_2) + \overline{a_3}c_1(a_3\overline{c_1} - a_2\overline{c_2}) - \overline{a_1}a_3c_1\overline{c_3} - a_1\overline{a_3c_1}c_3 + a_2\overline{a_2}c_1\overline{c_3} + a_1\overline{a_3}c_2\overline{c_2} \\
=&a_1\overline{c_3} + \overline{a_3}c_1 - a_1\overline{a_3}(c_1\overline{c_3} + \overline{c_1}c_3 - c_2\overline{c_2}) - c_1\overline{c_3}(a_1\overline{a_3} + \overline{a_1}a_3 - a_2\overline{a_2}) \\
=&a_1\overline{c_3} + \overline{a_3}c_1.
\end{aligned}$$

$$\begin{aligned}
&(a_2c_3 - a_3c_2)(\overline{a_1c_3} - \overline{a_3c_1}) \\
=&a_2\overline{c_3}(1 - \overline{a_3}c_1 + \overline{a_2}c_2) + \overline{a_3}c_2(1 + a_2\overline{c_2} - a_1\overline{c_3}) - \overline{a_1}a_3c_2\overline{c_3} - a_2\overline{a_3c_1}c_3 \\
=&a_2\overline{c_3} + \overline{a_3}c_2 - a_2\overline{a_3}(c_1\overline{c_3} + \overline{c_1}c_3 - c_2\overline{c_2}) + c_2\overline{c_3}(a_2\overline{a_2} - a_1\overline{a_3} - \overline{a_1}a_3) \\
=&a_2\overline{c_3} + \overline{a_3}c_2.
\end{aligned}$$

$$\begin{aligned}
&(a_2c_3 - a_3c_2)(\overline{a_2c_3} - \overline{a_3c_2}) \\
=&a_3\overline{c_3}(1 - \overline{a_3}c_1 - \overline{a_1}c_3) + \overline{a_3}c_3(1 - a_3\overline{c_1} - a_1\overline{c_3}) + a_2\overline{a_2}c_3\overline{c_3} + a_3\overline{a_3}c_2\overline{c_2} \\
=&a_3\overline{c_3} + \overline{a_3}c_3 - a_3\overline{a_3}(c_1\overline{c_3} + \overline{c_1}c_3 - c_2\overline{c_2}) - c_3\overline{c_3}(\overline{a_1}a_3 + a_1\overline{a_3} - a_2\overline{a_2}) \\
=&a_3\overline{c_3} + \overline{a_3}c_3.
\end{aligned}$$

By these identities, we get the desired formula.

Similarly, we have

$$F_{1,2} = \overline{v}|j(\gamma, Z)|^{-2}, \quad F_{2,1} = v|j(\gamma, Z)|^{-2}, \quad F_{2,2} = |j(\gamma, Z)|^{-2},$$

and

$$H_1 = -k|j(\gamma, Z)|^{-2}, \quad H_2 = 0, \quad G_1 = k|j(\gamma, Z)|^{-2}, \quad G_2 = 0.$$

Therefore,

$$\begin{aligned}
&L_k(f(\gamma(Z))j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k)\\
=&j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k(z_1+\overline{z_1}-z_2\overline{z_2})|j(\gamma,Z)|^{-2}\\
&\times\left[(u+\overline{u})\frac{\partial^2}{\partial u\partial\overline{u}}+\frac{\partial^2}{\partial v\partial\overline{v}}+v\frac{\partial^2}{\partial\overline{u}\partial v}+\overline{v}\frac{\partial^2}{\partial u\partial\overline{v}}-k\left(\frac{\partial}{\partial u}-\frac{\partial}{\partial\overline{u}}\right)\right]f(\gamma(Z))\\
=&j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k(L_kf)(\gamma(Z)).
\end{aligned}$$

□

For the Picard modular group $\Gamma = U(2,1;\mathcal{O})$, the Eisenstein series of weight k is defined by

$$E(Z;k,s)=\sum_{\gamma\in\Gamma\cap N\backslash\Gamma} j(\overline{\gamma},\overline{Z})^k j(\gamma,Z)^{-k}\rho(\gamma,Z)^s,\quad\text{for}\quad \mathrm{Re}(s)>2, k\in\mathbb{Z}. \tag{4.7}$$

THEOREM 4.3. *The Eisenstein series satisfies the following equation:*

$$L_kE(Z;k,s)=s(s-2)E(Z;k,s). \tag{4.8}$$

Proof. First, we note that

$$L_k\rho^s=L\rho^s+\rho\cdot k\left(\frac{\partial}{\partial z_1}-\frac{\partial}{\partial\overline{z_1}}\right)\rho^s=s(s-2)\rho^s.$$

By Proposition 4.2, we have

$$\begin{aligned}
L_kE(Z;k,s)&=\sum_{\gamma\in\Gamma\cap N\backslash\Gamma}L_k[\rho(\gamma(Z))^sj(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k]\\
&=\sum_{\gamma\in\Gamma\cap N\backslash\Gamma}j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k(L_k\rho^s)(\gamma(Z))\\
&=s(s-2)\sum_{\gamma\in\Gamma\cap N\backslash\Gamma}j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k\rho^s(\gamma(Z))\\
&=s(s-2)E(Z;k,s).
\end{aligned}$$

□

Set

$$J(\gamma,Z):=\frac{j(\gamma,Z)}{|j(\gamma,Z)|}. \tag{4.9}$$

Then $J(\gamma, Z) = e^{i \arg j(\gamma, Z)}$.

A generalized automorphic form on the Picard modular group $\Gamma = U(2,1;\mathcal{O})$ of weight $3k$ is given by

$$f(\gamma(Z)) = \frac{j(\gamma, Z)^{3k}}{|j(\gamma, Z)|^{3k}} f(Z) \quad \text{for} \quad \gamma \in \Gamma. \tag{4.10}$$

Put

$$H_k(Z, W) = \frac{\rho(Z, W)^{2k}}{|\rho(Z, W)|^{2k}} \quad \text{for} \quad Z, W \in \mathfrak{S}_2. \tag{4.11}$$

Definition 4.4. For any function $\Phi \in C^\infty(\mathbb{R})$, the associated real point-pair invariant is given by

$$k(Z, W) = \Phi\left[\frac{|\rho(Z, W)|^2}{\rho(Z)\rho(W)}\right], \tag{4.12}$$

where $Z, W \in \mathfrak{S}_2$.

PROPOSITION 4.5. *The following formulas hold:*

1. $k(Z, W) = k(g(Z), g(W))$*, for any* $g \in G$.
2. $k(Z, W) = k(W, Z)$.

Definition 4.6. Let $\Phi \in C^\infty(\mathbb{R})$ be a real-valued function. The complex point-pair invariant is given by

$$K_k(Z, W) = H_k(Z, W)\Phi\left[\frac{|\rho(Z, W)|^2}{\rho(Z)\rho(W)}\right], \tag{4.13}$$

where $Z, W \in \mathfrak{S}_2$.

PROPOSITION 4.7. *The following formulas hold:*

1. $H_k(g(Z), g(W)) = J(g, Z)^{2k} H_k(Z, W) J(g, W)^{-2k}$*, for* $g \in G$.
2. $K_k(g(W), g(W)) = J(g, Z)^{2k} K_k(Z, W) J(g, W)^{-2k}$*, for* $g \in G$.
3. $|H_k(Z, W)| = 1$*, and* $H_k(Z, W) = \overline{H_k(W, Z)}$.
4. $K_k(Z, W) = \overline{K_k(W, Z)}$.

Proof. Since $\rho(g(Z), g(W))\overline{j(g, Z)} j(g, W) = \rho(Z, W)$ for $g \in G$. We have

$$H_k(g(Z), g(W)) = \frac{|j(g, Z)|^{2k}}{\overline{j(g, Z)}^{2k}} H_k(Z, W) \frac{|j(g, W)|^{2k}}{j(g, W)^{2k}} = J(g, Z)^{2k} H_k(Z, W) J(g, W)^{-2k}.$$

The others are trivial.

□

Lemma 4.8. *The following formula holds:*

$$L_kK_k(Z,W) = H_k(Z,W)\left[(\sigma^2-\sigma)\Phi''(\sigma)+(3\sigma-1)\Phi'(\sigma)+\frac{k^2}{\sigma}\Phi(\sigma)\right], \tag{4.14}$$

where $\sigma(Z,W) = \frac{|\rho(Z,W)|^2}{\rho(Z)\rho(W)}$.

Proof. It is obtained by a straightforward calculation.

□

By $L_kK_k(Z,W) = s(s-2)K_k(Z,W)$, one has

$$\Phi''(\sigma)+\left(\frac{1}{\sigma}+\frac{2}{\sigma-1}\right)\Phi'(\sigma)+\frac{1}{\sigma(\sigma-1)}\left[\frac{k^2}{\sigma}-s(s-2)\right]\Phi(\sigma)=0. \tag{4.15}$$

It is known that the Fuchsian equation with three regular singularities a, b, c is given by

$$\begin{aligned}&\frac{d^2w}{dz^2}+\left[\frac{1-\alpha_1-\alpha_2}{z-a}+\frac{1-\beta_1-\beta_2}{z-b}+\frac{1-\gamma_1-\gamma_2}{z-c}\right]\frac{dw}{dz}+\frac{1}{(z-a)(z-b)(z-c)}\\ &\times\left[\frac{\alpha_1\alpha_2(a-b)(a-c)}{z-a}+\frac{\beta_1\beta_2(b-c)(b-a)}{z-b}+\frac{\gamma_1\gamma_2(c-a)(c-b)}{z-c}\right]w=0,\end{aligned}$$

where (α_1,α_2), (β_1,β_2), (γ_1,γ_2) are exponents belonging to a, b, c, respectively, and, they satisfy that $\alpha_1+\alpha_2+\beta_1+\beta_2+\gamma_1+\gamma_2=1$. The solution of this equation is given in Riemann P-notation by

$$w(z) = P\left\{\begin{matrix} a & b & c \\ \alpha_1 & \beta_1 & \gamma_1\,;z \\ \alpha_2 & \beta_2 & \gamma_2 \end{matrix}\right\}.$$

When $c=\infty$, it reduces to

$$\begin{aligned}&\frac{d^2w}{dz^2}+\left[\frac{1-\alpha_1-\alpha_2}{z-a}+\frac{1-\beta_1-\beta_2}{z-b}\right]\frac{dw}{dz}+\frac{1}{(z-a)(z-b)}\\ &\times\left[\frac{\alpha_1\alpha_2(a-b)}{z-a}+\frac{\beta_1\beta_2(b-a)}{z-b}+\gamma_1\gamma_2\right]w=0.\end{aligned}$$

Now, in our case,

$$a=0, b=1, c=\infty, \alpha_1=-|k|, \alpha_2=|k|, \beta_1=0, \beta_2=-1, \gamma_1=s, \gamma_2=2-s,$$

and

$$\begin{aligned}\Phi(\sigma) &= \Phi_{|k|}(\sigma)\\ &=P\left\{\begin{matrix} 0 & 1 & \infty & \\ -|k| & 0 & s & ;\sigma \\ |k| & -1 & 2-s & \end{matrix}\right\}\\ &=\sigma^{-|k|}P\left\{\begin{matrix} 0 & 1 & \infty & \\ 0 & 0 & s-|k| & ;\sigma \\ 2|k| & -1 & 2-s-|k| & \end{matrix}\right\}\\ &=\sigma^{-|k|}F(s-|k|,2-s-|k|;1-2|k|;\sigma)\\ &=\sigma^{-|k|}(1-\sigma)^{|k|-s}F(s-|k|,s-1-|k|;1-2|k|;\frac{\sigma}{\sigma-1})\\ &=\sigma^{-|k|}(1-\sigma)^{|k|-s}F(s-|k|,s-1-|k|;2s-1;-\frac{1}{\sigma-1}).\end{aligned}$$

Now, we set

$$\begin{aligned}K(Z,W;k,s) =&\frac{\pi^{\frac{3}{2}}\Gamma(\frac{3}{2}-s)}{(|k|+1-s)\Gamma(2-s)}(\frac{1}{4})^{s-1}\rho(Z,W)^k\rho(\overline{Z},\overline{W})^{-k}\\ &\times \sigma^{-|k|}(\sigma-1)^{|k|-s}F(s-|k|,s-1-|k|;2s-1;-\frac{1}{\sigma-1}).\end{aligned} \tag{4.16}$$

It has the following property:

$$K(\gamma(Z),\gamma(Z');k,s) = j(\gamma,Z)^k j(\overline{\gamma},\overline{Z})^{-k} j(\gamma,Z')^{-k} j(\overline{\gamma},\overline{Z'})^k K(Z,Z';k,s). \tag{4.17}$$

Moreover, note that $\rho(Z',Z)=\rho(\overline{Z},\overline{Z'})$, we have

$$K(Z',Z;k,s) = K(Z,Z';-k,s).$$

The automorphic Green function of weight k is given by

$$G(Z,Z';k,s) = \sum_{\gamma\in\Gamma} K(Z,\gamma(Z');k,s), \quad \text{for} \quad \mathrm{Re}(s) > \delta(\Gamma). \tag{4.18}$$

By the same method, we can prove that the right hand side of (4.18) is convergent if $\mathrm{Re}(s)>\delta(\Gamma)$.

The automorphic Green function of weight k has the following properties:

$$G(Z,Z';k,s) = G(Z',Z;-k,s), \quad L_k(Z)G(Z,Z';k,s) = s(s-2)G(Z,Z';k,s),$$

Furthermore,

$$\begin{aligned}L_{-k}(Z')G(Z,Z';k,s) &= L_{-k}(Z')G(Z',Z;-k,s)\\ &=s(s-2)G(Z',Z;-k,s) = s(s-2)G(Z,Z';k,s).\end{aligned}$$

The Poisson kernel of weight k is given by

$$P_k(Z,W;s)=\rho(Z)^s\rho(Z,W)^{k-s}\rho(\overline{Z},\overline{W})^{-k-s}, \tag{4.19}$$

where $Z\in\mathfrak{S}_2$, $W\in\partial\mathfrak{S}_2$. By

$$\rho(\gamma(Z),\gamma(W))\overline{j(\gamma,Z)}j(\gamma,W)=\rho(Z,W),\quad j(\gamma,\gamma^{-1}(W))=j(\gamma^{-1},W)^{-1},$$

we have

$$P_k(\gamma(Z),W;s)=P_k(Z,\gamma^{-1}(W);s)j(\gamma,Z)^kj(\overline{\gamma},\overline{Z})^{-k}j(\gamma^{-1},W)^{k-s}j(\overline{\gamma}^{-1},\overline{W})^{-k-s}.$$

The Eisenstein series with respect to $\partial\mathfrak{S}_2$ of weight k is defined by

$$E(Z,W;k,s)=\sum_{\gamma\in\Gamma}j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^kP_k(\gamma(Z),W;s),\quad\text{for}\quad\mathrm{Re}(s)>\delta(\Gamma). \tag{4.20}$$

Here Γ is a discrete subgroup of $U(2,1)$. Now, we have the following formula:

$$\lim_{\rho(Z)\to 0}\rho(Z)^{-s}K_k(Z,Z')=\frac{\pi^{\frac{3}{2}}\Gamma(\frac{3}{2}-s)}{(|k|+1-s)\Gamma(2-s)}(\frac{1}{4})^{s-1}P_k(Z',W;s),$$

where $Z\to W\in\partial\mathfrak{S}_2$ as $\rho(Z)\to 0$. Consequently,

$$\lim_{\rho(Z)\to 0}\rho(Z)^{-s}G(Z,Z';k,s)=\frac{\pi^{\frac{3}{2}}\Gamma(\frac{3}{2}-s)}{(|k|+1-s)\Gamma(2-s)}(\frac{1}{4})^{s-1}E(Z',W;k,s).$$

LEMMA 4.9. *The Eisenstein series of weight k satisfies the following three equations:*

(1) $L_kE(Z,W;k,s)=s(s-2)E(Z,W;k,s)$.
(2) $E(\gamma(Z),W;k,s)=E(Z,W;k,s)$, *for* $\gamma\in\Gamma$.
(3) $E(Z,g(W);k,s)=j(g,W)^{s-k}j(\overline{g},\overline{W})^{s+k}E(Z,W;k,s)$, *where* $g\in\Gamma$.

Proof. In fact,

$$L_kP_k(Z,W;s)=s(s-2)P_k(Z,W;s)+(k^2-s^2)\rho(W)P_k(Z,W;s+1).$$

Since $\rho(W)=0$, $L_kP_k(Z,W;s)=s(s-2)P_k(Z,W;s)$. By Proposition 4.2,

$$\begin{aligned}L_kE(Z,W;k,s)=&\sum_{\gamma\in\Gamma}L_k[P_k(\gamma(Z),W;s)j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k]\\=&\sum_{\gamma\in\Gamma}j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^k(L_kP_k)(\gamma(Z),W;s)\\=&s(s-2)\sum_{\gamma\in\Gamma}j(\gamma,Z)^{-k}j(\overline{\gamma},\overline{Z})^kP_k(\gamma(Z),W;s)\\=&s(s-2)E(Z,W;k,s).\end{aligned}$$

To prove the third identity, we note that

$$P_k(\gamma(Z),\gamma(W);s)=j(\gamma,Z)^k j(\overline{\gamma},\overline{Z})^{-k} j(\gamma,W)^{s-k} j(\overline{\gamma},\overline{W})^{s+k} P_k(Z,W;s).$$

By Proposition 2.9, we have

$$\begin{aligned}E(Z,g(W);k,s)=&\sum_{\gamma\in\Gamma} j(\gamma,Z)^{-k} j(\overline{\gamma},\overline{Z})^k P_k(\gamma(Z),g(W);s)\\ =&\sum_{\gamma\in\Gamma} j(g,\gamma(Z))^{-k} j(\overline{g},\overline{\gamma}(\overline{Z}))^k j(\gamma,Z)^{-k} j(\overline{\gamma},\overline{Z})^k P_k(g(\gamma(Z)),g(W);s)\\ =&\sum_{\gamma\in\Gamma} j(\gamma,Z)^{-k} j(\overline{\gamma},\overline{Z})^k P_k(\gamma(Z),W;s) j(g,W)^{s-k} j(\overline{g},\overline{W})^{s+k}\\ =&j(g,W)^{s-k} j(\overline{g},\overline{W})^{s+k} E(Z,W;k,s).\end{aligned}$$

□

THEOREM 4.10. *Set*

$$\phi(Z,Z';k,s)=\int_{\partial\mathfrak{S}_2} P_k(Z,W;s)P_{-k}(Z',W;2-s)dm(W). \tag{4.21}$$

Then the following formula holds:

$$\begin{aligned}\phi(Z,Z';k,s)&=K(Z,Z';k,s)+K(Z,Z';k,2-s)\\ =\phi(Z',Z;-k,2-s)&=K(Z',Z;-k,s)+K(Z',Z;-k,2-s),\end{aligned} \tag{4.22}$$

where $k=0$ or $k=\pm1$.

Proof. In fact, if $k=0$, this is Theorem 3.7. Now, we only consider the nontrivial case $k=\pm1$.

For $\gamma\in G$,

$$\begin{aligned}\phi(\gamma(Z),\gamma(Z');k,s)=&\int_{\partial\mathfrak{S}_2} P_k(\gamma(Z),W;s)P_{-k}(\gamma(Z'),W;2-s)dm(W)\\ =&j(\gamma,Z)^k j(\overline{\gamma},\overline{Z})^{-k} j(\gamma,Z')^{-k} j(\overline{\gamma},\overline{Z'})^k \int_{\partial\mathfrak{S}_2} P_k(Z,\gamma^{-1}(W);s)\\ &\times P_{-k}(Z',\gamma^{-1}(W);2-s) j(\gamma^{-1},W)^{-2} j(\overline{\gamma}^{-1},\overline{W})^{-2} dm(W).\end{aligned}$$

Since $dm(\gamma^{-1}(W))=|j(\gamma^{-1},W)|^{-4}dm(W)$, we have

$$\phi(\gamma(Z),\gamma(Z');k,s)=j(\gamma,Z)^k j(\overline{\gamma},\overline{Z})^{-k} j(\gamma,Z')^{-k} j(\overline{\gamma},\overline{Z'})^k \phi(Z,Z';k,s). \tag{4.23}$$

Thus, $\phi(Z,Z';k,s)$ is a point-pair invariant covariant with respect to the weight k.

By the double transitivity of $U(2,1)$ on $\mathfrak{S}_2$, there exists a $\gamma \in U(2,1)$, such that

$$Z = \gamma(\frac{1}{2}\rho, 0), \quad Z' = \gamma(\frac{1}{2}\rho', 0).$$

Let ρ and ρ' stand for $(\rho, 0)$ and $(\rho', 0)$, respectively. Set $\gamma = \begin{pmatrix} * & * & * \\ * & * & * \\ a_1 & a_2 & a_3 \end{pmatrix}$, then

$$|j(\gamma, \frac{1}{2}\rho)|^2 = |\frac{1}{2}\rho a_1 + a_3|^2 = \frac{1}{4}\rho^2|a_1|^2 + \frac{1}{2}\rho|a_2|^2 + |a_3|^2.$$

Denote

$$\mu = \frac{\frac{1}{2}(\rho + \rho')}{\frac{1}{4}\rho^2|a_1|^2 + \frac{1}{2}\rho|a_2|^2 + |a_3|^2}, \tag{4.24}$$

then $\mu > 0$.

Since

$$\begin{aligned} \phi(Z, Z'; k, s) &= \phi(\gamma(\frac{1}{2}\rho), \gamma(\frac{1}{2}\rho'); k, s) \\ &= j(\gamma, \frac{1}{2}\rho)^k j(\overline{\gamma}, \frac{1}{2}\rho)^{-k} j(\gamma, \frac{1}{2}\rho')^{-k} j(\overline{\gamma}, \frac{1}{2}\rho')^k \phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s). \end{aligned}$$

By (4.24), we have

$$\rho(Z, Z') = \rho(\gamma(\frac{1}{2}\rho), \gamma(\frac{1}{2}\rho')) = \frac{\frac{1}{2}(\rho + \rho')}{j(\overline{\gamma}, \frac{1}{2}\rho) j(\gamma, \frac{1}{2}\rho')} = \mu \frac{j(\gamma, \frac{1}{2}\rho)}{j(\gamma, \frac{1}{2}\rho')}.$$

Hence,

$$\begin{aligned} \phi(Z, Z'; k, s) &= [\mu^{-1}\rho(Z, Z')]^k [\mu^{-1}\rho(\overline{Z}, \overline{Z'})]^{-k} \phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) \\ &= \rho(Z, Z')^k \rho(\overline{Z}, \overline{Z'})^{-k} \phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s). \end{aligned} \tag{4.25}$$

For $W \in \partial\mathfrak{S}_2$, i.e., $w_1 = \frac{|w|^2}{2} + iv$, $w_2 = w$ with $w \in \mathbb{C}$, $v \in \mathbb{R}$.

$$\rho(Z, W) = \frac{\rho + |w|^2}{2} + iv, \quad \rho(Z', W) = \frac{\rho' + |w|^2}{2} + iv.$$

Therefore,

$$\begin{aligned} \phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) &= 16\rho^s {\rho'}^{2-s} \int_{\mathbb{R}} \int_{\mathbb{C}} \frac{1}{|\rho + |w|^2 + 2iv|^{2s}} \frac{1}{|\rho' + |w|^2 + 2iv|^{4-2s}} \\ &\quad \times \left(\frac{\rho + |w|^2 + 2iv}{\rho + |w|^2 - 2iv}\right)^k \left(\frac{\rho' + |w|^2 + 2iv}{\rho' + |w|^2 - 2iv}\right)^{-k} dw d\overline{w} dv. \end{aligned}$$

Set $w = re^{i\theta}$ and $u = r^2$, then

$$\begin{aligned}\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) =&16\pi\rho^s{\rho'}^{2-s}\int_{-\infty}^{\infty}\int_0^{\infty}\frac{1}{|\rho+u+2iv|^{2s}}\frac{1}{|\rho'+u+2iv|^{4-2s}}\\ &\times\left(\frac{\rho+u+2iv}{\rho+u-2iv}\right)^k\left(\frac{\rho'+u+2iv}{\rho'+u-2iv}\right)^{-k}dudv\\ =&8\pi\rho^s{\rho'}^{2-s}\int_{-\infty}^{\infty}\int_0^{\infty}\frac{1}{|\rho+u+iv|^{2s}}\frac{1}{|\rho'+u+iv|^{2(2-s)}}\\ &\times\exp(2ki(\arctan\frac{v}{\rho+u}-\arctan\frac{v}{\rho'+u}))dudv.\end{aligned}$$

By

$$\arctan\frac{v}{\rho+u}-\arctan\frac{v}{\rho'+u}=\arctan\frac{(\rho'-\rho)v}{(\rho+u)(\rho'+u)+v^2},$$

we have

$$\begin{aligned}\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) =&8\pi\rho^s{\rho'}^{2-s}\int_{-\infty}^{\infty}\int_0^{\infty}\frac{1}{[(\rho+u)^2+v^2]^s}\frac{1}{[(\rho'+u)^2+v^2]^{2-s}}\\ &\times\exp(2ki\arctan\frac{(\rho'-\rho)v}{(\rho+u)(\rho'+u)+v^2})dudv\\ =&16\pi\rho^s{\rho'}^{2-s}\int_0^{\infty}\int_0^{\infty}\frac{1}{[(\rho+u)^2+v^2]^s}\frac{1}{[(\rho'+u)^2+v^2]^{2-s}}\\ &\times\cos 2k\arctan\frac{(\rho'-\rho)v}{(\rho+u)(\rho'+u)+v^2}dudv.\end{aligned}$$

Assume that $\rho < \rho'$, set $\lambda = \frac{\rho}{\rho'}$, $v = (\rho'-\rho)w$ and $l = \frac{\rho+u}{\rho'+u}$, then $u = \frac{l\rho'-\rho}{1-l}$, $du = \frac{\rho'-\rho}{(1-l)^2}dl$. We have

$$\begin{aligned}\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) =&16\pi\lambda^s(1-\lambda)^{-2}\int_0^{\infty}\int_{\lambda}^1\left[w^2+\frac{l^2}{(1-l)^2}\right]^{-s}\left[w^2+\frac{1}{(1-l)^2}\right]^{s-2}\\ &\times\cos 2k\arctan\left(\frac{w}{w^2+\frac{l}{(1-l)^2}}\right)(1-l)^{-2}dldw.\end{aligned}$$

Set $w = \frac{\sqrt{v}}{1-l}$, then

$$\begin{aligned}\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) =&8\pi\lambda^s(1-\lambda)^{-2}\int_{\lambda}^1(1-l)dl\int_0^{\infty}v^{-\frac{1}{2}}(1+v)^{s-2}(l^2+v)^{-s}\\ &\times\cos 2k\arctan\left[\frac{(1-l)\sqrt{v}}{v+l}\right]dv.\end{aligned}$$

Let $x = \arctan \frac{(1-l)\sqrt{v}}{v+l}$, then

$$\cos 2x = \frac{1-\tan^2 x}{1+\tan^2 x} = \frac{(v+l)^2-(1-l)^2 v}{(v+l)^2+(1-l)^2 v}.$$

We have

$$\cos 2kx = \cos\left(k \arccos \frac{(v+l)^2-(1-l)^2 v}{(v+l)^2+(1-l)^2 v}\right).$$

It is known that the Tchebichef polynomial

$$T_n(x) = \cos(n \arccos x) = F(-n, n; \frac{1}{2}; \frac{1-x}{2}).$$

Hence

$$\cos 2kx = F(-k, k; \frac{1}{2}; \frac{(1-l)^2 v}{v^2+(1+l^2)v+l^2}).$$

Therefore,

$$\begin{aligned}\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) =& 8\pi\lambda^s(1-\lambda)^{-2}\int_\lambda^1 (1-l)dl \int_0^\infty v^{-\frac{1}{2}}(1+v)^{s-2}(l^2+v)^{-s} \\ &\times F(-k, k; \frac{1}{2}; \frac{(1-l)^2 v}{(1+v)(l^2+v)})dv.\end{aligned}$$

We know that

$$F(-k, k; \frac{1}{2}; x) = \sum_{n=0}^{|k|} \frac{(-k)_n (k)_n}{n!(\frac{1}{2})_n} x^n.$$

Thus

$$\begin{aligned}\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) =& 8\pi\lambda^s(1-\lambda)^{-2}\sum_{n=0}^{|k|} \frac{(-k)_n (k)_n}{n!(\frac{1}{2})_n} \int_\lambda^1 (1-l)^{2n+1} l^{-2(s+n)} dl \\ &\times \int_0^\infty v^{n-\frac{1}{2}}(1+v)^{s-n-2}(1+vl^{-2})^{-s-n} dv.\end{aligned}$$

By the following formula

$$F(a, b; c; 1-z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^\infty s^{b-1}(1+s)^{a-c}(1+sz)^{-a} ds$$

for $\mathrm{Re}(c) > \mathrm{Re}(b) > 0$, $|\arg(z)| < \pi$, we have

$$\begin{aligned}&\int_0^\infty v^{n-\frac{1}{2}}(1+v)^{s-n-2}(1+vl^{-2})^{-s-n} dv \\ =& \frac{\Gamma(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{\Gamma(2n+2)} F(s+n, \frac{1}{2}+n; 2n+2; 1-l^{-2}).\end{aligned}$$

Hence,

$$\begin{aligned}\phi(\frac{1}{2}\rho,\frac{1}{2}\rho';k,s)=&8\pi\lambda^s(1-\lambda)^{-2}\sum_{n=0}^{|k|}\frac{(-k)_n(k)_n}{n!(\frac{1}{2})_n}\frac{\Gamma(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{\Gamma(2n+2)}\\&\times\int_\lambda^1(1-l)^{2n+1}l^{-2(s+n)}F(s+n,\frac{1}{2}+n;2n+2;1-l^{-2})dl.\end{aligned}$$

Note that

$$\frac{1}{n!(\frac{1}{2})_n}\frac{\Gamma(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{\Gamma(2n+2)}=\frac{\pi}{2^{2n+1}(n!)^2}.$$

We have

$$\begin{aligned}\phi(\frac{1}{2}\rho,\frac{1}{2}\rho';k,s)=&4\pi^2\lambda^s(1-\lambda)^{-2}\sum_{n=0}^{|k|}\frac{(-k)_n(k)_n}{2^{2n}(n!)^2}\int_\lambda^1(1-l)^{2n+1}l^{-2(s+n)}\\&\times F(s+n,\frac{1}{2}+n;2n+2;1-l^{-2})dl.\end{aligned}$$

(1) $z=1-l^{-2}$, $\frac{z}{z-1}=1-l^2$.
By $F(a,b;c;z)=(1-z)^{-a}F(a,c-b;c;\frac{z}{z-1})$, we have

$$F(s+n,\frac{1}{2}+n;2n+2;1-l^{-2})=l^{2(s+n)}F(s+n,n+\frac{3}{2};2n+2;1-l^2).$$

(2) $z=1-l^2$, $1-z=l^2$.
By

$$\begin{aligned}F(a,b;c;z)=&\frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}F(a,b;a+b-c+1;1-z)\\&+\frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)}(1-z)^{c-a-b}F(c-a,c-b;c-a-b+1;1-z),\end{aligned}$$

where $|\arg(1-z)|<\pi$, we have

$$\begin{aligned}&F(s+n,n+\frac{3}{2};2n+2;1-l^2)\\=&\frac{\Gamma(2n+2)\Gamma(\frac{1}{2}-s)}{\Gamma(n-s+2)\Gamma(n+\frac{1}{2})}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)\\&+\frac{\Gamma(2n+2)\Gamma(s-\frac{1}{2})}{\Gamma(s+n)\Gamma(n+\frac{3}{2})}l^{1-2s}F(n-s+2,n+\frac{1}{2};\frac{3}{2}-s;l^2).\end{aligned}$$

Therefore,

$$\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) = 4\pi^2 \lambda^s (1-\lambda)^{-2} \sum_{n=0}^{|k|} \frac{(-k)_n (k)_n}{2^{2n}(n!)^2}(F_1 + F_2),$$

where

$$F_1 = \frac{(2n+1)!\Gamma(\frac{1}{2}-s)}{\Gamma(n-s+2)\Gamma(n+\frac{1}{2})} \int_\lambda^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2)dl,$$

$$F_2 = \frac{(2n+1)!\Gamma(s-\frac{1}{2})}{\Gamma(s+n)\Gamma(n+\frac{3}{2})} \int_\lambda^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2)dl.$$

It is known that the incomplete beta function is defined by (see [Er], p.87)

$$B_x(p,q) = \int_0^x t^{p-1}(1-t)^{q-1}dt.$$

Set $I_x(p,q) = \frac{B_x(p,q)}{B_1(p,q)}$. Note that $B_1(p,q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$.

Hence,

$$\int_\lambda^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2)dl$$
$$= \sum_{m=0}^{\infty} \frac{(s+n)_m (n+\frac{3}{2})_m}{m!(s+\frac{1}{2})_m} [B_1(2m+1, 2n+2) - B_\lambda(2m+1, 2n+2)].$$

Similarly,

$$\int_\lambda^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2)dl$$
$$= \sum_{m=0}^{\infty} \frac{(2-s+n)_m (n+\frac{1}{2})_m}{m!(\frac{3}{2}-s)_m} [B_1(2m-2s+2, 2n+2) - B_\lambda(2m-2s+2, 2n+2)].$$

We have

$$\begin{aligned}(4.26)\qquad &\sum_{m=0}^{\infty} \frac{(s+n)_m (n+\frac{3}{2})_m}{m!(s+\frac{1}{2})_m} B_1(2m+1, 2n+2) = \sum_{m=0}^{\infty} \frac{(s+n)_m}{(s+\frac{1}{2})_m} \frac{(2m-1)!!(2n)!!}{(2m+2n+2)!!} \\ &= \frac{1}{2n+2} \sum_{m=0}^{\infty} \frac{(s+n)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (n+2)_m}.\end{aligned}$$

By

$$(n+2)_m = \frac{(n+m+1)!}{(n+1)!}, \quad (\frac{1}{2})_m = \frac{(-\frac{1}{2}-n)_{m+n+1}}{(-\frac{1}{2}-n)_{n+1}},$$

$$(s+\frac{1}{2})_m = \frac{(s-n-\frac{1}{2})_{m+n+1}}{(s-n-\frac{1}{2})_{n+1}}, \quad (s+n)_m = \frac{(s-1)_{m+n+1}}{(s-1)_{n+1}},$$

(4.26) is equal to

$$\begin{aligned}&\frac{1}{2n+2}\frac{(n+1)!(s-n-\frac{1}{2})_{n+1}}{(s-1)_{n+1}(-\frac{1}{2}-n)_{n+1}}\sum_{m=0}^{\infty}\frac{(s-1)_{m+n+1}(-\frac{1}{2}-n)_{m+n+1}}{(s-n-\frac{1}{2})_{m+n+1}(m+n+1)!}\\ =&\frac{1}{2n+2}\frac{(n+1)!(s-n-\frac{1}{2})_{n+1}}{(s-1)_{n+1}(-\frac{1}{2}-n)_{n+1}}[F(s-1,-\frac{1}{2}-n;s-n-\frac{1}{2};1)\\ &-\sum_{r=0}^{n}\frac{(s-1)_r(-\frac{1}{2}-n)_r}{(s-n-\frac{1}{2})_r r!}].\end{aligned} \tag{4.27}$$

Similarly,

$$\begin{aligned}&\sum_{m=0}^{\infty}\frac{(2-s+n)_m(n+\frac{1}{2})_m}{m!(\frac{3}{2}-s)_m}B_1(2m-2s+2,2n+2)\\ =&\frac{(2n+1)!\Gamma(2-2s)}{\Gamma(4-2s+2n)}\sum_{m=0}^{\infty}\frac{(1-s)_m(n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!}\\ =&\frac{(2n+1)!\Gamma(2-2s)}{\Gamma(4-2s+2n)}F(1-s,n+\frac{1}{2};\frac{5}{2}-s+n;1).\end{aligned} \tag{4.28}$$

Therefore,

$$\begin{aligned}&\sum_{m=0}^{\infty}\frac{(s+n)_m(n+\frac{3}{2})_m}{m!(s+\frac{1}{2})_m}B_\lambda(2m+1,2n+2)\\ =&\sum_{m=0}^{\infty}\frac{(s+n)_m(n+\frac{3}{2})_m}{m!(s+\frac{1}{2})_m}B_1(2m+1,2n+2)I_\lambda(2m+1,2n+2)\\ =&\frac{1}{2n+2}\sum_{m=0}^{\infty}\frac{(s+n)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(n+2)_m}I_\lambda(2m+1,2n+2).\end{aligned}$$

Now, we need the following lemma:

LEMMA 4.11. *The explicit expression of $I_x(p,q)$: For $p \in \mathbb{C}$, $Re(p) > 0$ and $q \in \mathbb{Z}$ with $q > 0$, the following formula holds:*

$$I_x(p,q+1) = (p)_{q+1}\sum_{r=0}^{q}\frac{(-1)^r}{r!(q-r)!}\frac{x^{p+r}}{p+r}. \tag{4.29}$$

Proof.

$$\begin{aligned}I_x(p,q+1) &= \frac{(p)_{q+1}}{q!}B(p,q+1)I_x(p,q+1)\\ =&\frac{(p)_{q+1}}{q!}B_x(p,q+1) = \frac{(p)_{q+1}}{q!}\int_0^x t^{p-1}(1-t)^q dt\\ =&\frac{(p)_{q+1}}{q!}\sum_{r=0}^{q}(-1)^r C_q^r \frac{x^{p+r}}{p+r} = (p)_{q+1}\sum_{r=0}^{q}\frac{(-1)^r}{r!(q-r)!}\frac{x^{p+r}}{p+r}.\end{aligned}$$

□

By Lemma 4.11, we have

$$\begin{aligned}&\frac{1}{2n+2}\sum_{m=0}^{\infty}\frac{(s+n)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(n+2)_m}I_\lambda(2m+1,2n+2)\\ =&\frac{1}{2n+2}\sum_{m=0}^{\infty}\frac{(s+n)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(n+2)_m}(2m+1)_{2n+2}\sum_{r=0}^{2n+1}\frac{(-1)^r}{r!(2n+1-r)!}\frac{\lambda^{2m+r+1}}{2m+r+1}\\ =&\frac{1}{2n+2}\sum_{r=0}^{2n+1}\frac{(-1)^r}{r!(2n+1-r)!}\sum_{m=0}^{\infty}\frac{(s+n)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(n+2)_m}\\ &\times\frac{(2m+1)(2m+2)\cdots(2m+2n+2)}{2m+r+1}\lambda^{2m+r+1}.\end{aligned}$$

If $n=0$, then $r=0,1$.

(1) $r=0$,

$$\sum_{m=0}^{\infty}\frac{(s)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(2)_m}(2m+2)\lambda^{2m+1} = 2\lambda F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2).$$

(2) $r=1$,

$$\sum_{m=0}^{\infty}\frac{(s)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(2)_m}(2m+1)\lambda^{2m+2} = \frac{2s-1}{s-1}[F(s-1,\frac{1}{2};s-\frac{1}{2};\lambda^2)-1].$$

On the other hand, when $n=0$, (4.27) is equal to

$$\frac{2s-1}{2s-2}[1-F(s-1,-\frac{1}{2};s-\frac{1}{2};1)].$$

Hence,

$$\begin{aligned}&\int_\lambda^1(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=0}\\ =&-\lambda F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)+\frac{2s-1}{2s-2}F(s-1,\frac{1}{2};s-\frac{1}{2};\lambda^2)-\frac{2s-1}{2s-2}F(s-1,-\frac{1}{2};s-\frac{1}{2};1).\end{aligned}$$

If $n=1$, then $r=0,1,2,3$.

(1) $r=0$,

$$\begin{aligned}&\sum_{m=0}^{\infty}\frac{(s+1)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(3)_m}(2m+2)(2m+3)(2m+4)\lambda^{2m+1}\\=&8\lambda F(s+1,\frac{3}{2};s+\frac{1}{2};\lambda^2)+16\lambda F(s+1,\frac{1}{2};s+\frac{1}{2};\lambda^2).\end{aligned}$$

(2) $r=1$,

$$\sum_{m=0}^{\infty}\frac{(s+1)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(3)_m}(2m+1)(2m+3)(2m+4)\lambda^{2m+2}=8\frac{s-\frac{1}{2}}{s}[F(s,\frac{3}{2};s-\frac{1}{2};\lambda^2)-1].$$

(3) $r=2$,

$$\sum_{m=0}^{\infty}\frac{(s+1)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(3)_m}(2m+1)(2m+2)(2m+4)\lambda^{2m+3}=8\lambda^3F(s+1,\frac{3}{2};s+\frac{1}{2};\lambda^2).$$

(4) $r=3$,

$$\begin{aligned}&\sum_{m=0}^{\infty}\frac{(s+1)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(3)_m}(2m+1)(2m+2)(2m+3)\lambda^{2m+4}\\=&8\frac{s-\frac{1}{2}}{s}\lambda^2[F(s,\frac{3}{2};s-\frac{1}{2};\lambda^2)-1]-16\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}[F(s-1,\frac{1}{2};s-\frac{3}{2};\lambda^2)\\&-\frac{s-1}{2s-3}\lambda^2-1].\end{aligned}$$

On the other hand, when $n=1$, (4.27) is equal to

$$\frac{2}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}F(s-1,-\frac{3}{2};s-\frac{3}{2};1)-\frac{2}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}+\frac{s-\frac{1}{2}}{s}.$$

Therefore,

$$\begin{aligned}&\int_{\lambda}^{1}(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=1}\\=&\frac{2}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}F(s-1,-\frac{3}{2};s-\frac{3}{2};1)-\frac{2}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}F(s-1,\frac{1}{2};s-\frac{3}{2};\lambda^2)\\&-\frac{2}{3}\lambda F(s+1,\frac{1}{2};s+\frac{1}{2};\lambda^2)+(\frac{1}{3}\lambda^2+1)\frac{s-\frac{1}{2}}{s}F(s,\frac{3}{2};s-\frac{1}{2};\lambda^2)\\&-(\lambda^3+\frac{1}{3}\lambda)F(s+1,\frac{3}{2};s+\frac{1}{2};\lambda^2).\end{aligned}$$

Now, we need the following relation of Gauss between contiguous functions (see [Er], p.103, (36)):

$$(c-a-b)F-(c-a)F(a-1)+b(1-z)F(b+1)=0,$$

where F denotes $F(a,b;c;z)$ and $F(a\pm 1)$, $F(b\pm 1)$, and $F(c\pm 1)$ stands for $F(a\pm 1,b;c;z)$, $F(a,b\pm 1;c;z)$, and $F(a,b;c\pm 1;z)$ respectively.

We have

$$F(s-1,\frac{1}{2};s-\frac{3}{2};\lambda^2)=\frac{1}{2}F(s-2,\frac{1}{2};s-\frac{3}{2};\lambda^2)+\frac{1}{2}(1-\lambda^2)F(s-1,\frac{3}{2};s-\frac{3}{2};\lambda^2),$$

$$F(s+1,\frac{1}{2};s+\frac{1}{2};\lambda^2)=\frac{1}{2}F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)+\frac{1}{2}(1-\lambda^2)F(s+1,\frac{3}{2};s+\frac{1}{2};\lambda^2).$$

By [Er], p.104, (46),

$$F(a,b;c;1)=\frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)},\quad c\neq 0,-1,-2,\cdots,\mathrm{Re}(c)>\mathrm{Re}(a+b),$$

we have

$$F(s-1,-\frac{3}{2};s-\frac{3}{2};1)=\frac{\Gamma(s-\frac{3}{2})}{-2\sqrt{\pi}\Gamma(s)},\quad F(s-1,-\frac{1}{2};s-\frac{1}{2};1)=\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}.$$

Therefore,

$$\begin{aligned}&\int_\lambda^1(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=0}\\ =&-\lambda F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)+\frac{s-\frac{1}{2}}{s-1}F(s-1,\frac{1}{2};s-\frac{1}{2};\lambda^2)-\frac{1}{\sqrt{\pi}}\frac{\Gamma(s+\frac{1}{2})}{(s-1)\Gamma(s)},\end{aligned}$$

and

$$\begin{aligned}&\int_\lambda^1(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=1}\\ =&-\frac{1}{3}\frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}(s-1)\Gamma(s+1)}-\frac{1}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}F(s-2,\frac{1}{2};s-\frac{3}{2};\lambda^2)\\ &-\frac{1}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}(1-\lambda^2)F(s-1,\frac{3}{2};s-\frac{3}{2};\lambda^2)+(\frac{1}{3}\lambda^2+1)\frac{s-\frac{1}{2}}{s}F(s,\frac{3}{2};s-\frac{1}{2};\lambda^2)\\ &-\frac{1}{3}\lambda F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)-\frac{2}{3}\lambda(\lambda^2+1)F(s+1,\frac{3}{2};s+\frac{1}{2};\lambda^2).\end{aligned}$$

By [Er], p.111, (5),

$$F(a,b;2b;\frac{4z}{(1+z)^2})=(1+z)^{2a}F(a,a+\frac{1}{2}-b;b+\frac{1}{2};z^2),$$

we have

$$\begin{aligned}
&F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)=(1-\lambda)^{-2s}F(s,s;2s;-u^{-1}),\\
&F(s-1,\frac{1}{2};s-\frac{1}{2};\lambda^2)=(1-\lambda)^{-2(s-1)}F(s-1,s-1;2s-2;-u^{-1}),\\
&F(s-2,\frac{1}{2};s-\frac{3}{2};\lambda^2)=(1-\lambda)^{-2(s-2)}F(s-2,s-2;2s-4;-u^{-1}),\\
&F(s-1,\frac{3}{2};s-\frac{3}{2};\lambda^2)=(1-\lambda)^{-2(s-1)}F(s-1,s-2;2s-4;-u^{-1}),\\
&F(s,\frac{3}{2};s-\frac{1}{2};\lambda^2)=(1-\lambda)^{-2s}F(s,s-1;2s-2;-u^{-1}),\\
&F(s,\frac{1}{2};s+\frac{1}{2};\lambda^2)=(1-\lambda)^{-2s}F(s,s;2s;-u^{-1}),\\
&F(s+1,\frac{3}{2};s+\frac{1}{2};\lambda^2)=(1-\lambda)^{-2(s+1)}F(s+1,s;2s;-u^{-1}).
\end{aligned}$$

where $u=\frac{(\lambda-1)^2}{4\lambda}$. Hence,

$$\begin{aligned}
&\int_\lambda^1(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=0}=-\lambda(1-\lambda)^{-2s}F(s,s;2s;-u^{-1})\\
&+\frac{s-\frac{1}{2}}{s-1}(1-\lambda)^{-2(s-1)}F(s-1,s-1;2s-2;-u^{-1})-\frac{1}{\sqrt{\pi}}\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\Gamma(s)}.
\end{aligned}$$

$$\begin{aligned}
&\int_\lambda^1(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=1}=-\frac{1}{3}\frac{1}{\sqrt{\pi}}\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\Gamma(s+1)}\\
&-\frac{1}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}(1-\lambda)^{-2(s-2)}F(s-2,s-2;2s-4;-u^{-1})\\
&-\frac{1}{3}\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)}(1-\lambda^2)(1-\lambda)^{-2(s-1)}F(s-1,s-2;2s-4;-u^{-1})\\
&+(\frac{1}{3}\lambda^2+1)\frac{s-\frac{1}{2}}{s}(1-\lambda)^{-2s}F(s,s-1;2s-2;-u^{-1})\\
&-\frac{1}{3}\lambda(1-\lambda)^{-2s}F(s,s;2s;-u^{-1})-\frac{2}{3}\lambda(\lambda^2+1)(1-\lambda)^{-2(s+1)}F(s+1,s;2s;-u^{-1}).
\end{aligned}$$

Similarly,

$$\begin{aligned}
&\sum_{m=0}^{\infty} \frac{(2-s+n)_m(n+\frac{1}{2})_m}{m!(\frac{3}{2}-s)_m} B_\lambda(2m+2-2s, 2n+2)\\
=&\sum_{m=0}^{\infty} \frac{(2-s+n)_m(n+\frac{1}{2})_m}{m!(\frac{3}{2}-s)_m} B_1(2m+2-2s, 2n+2) I_\lambda(2m+2-2s, 2n+2)\\
=&\frac{(2n+1)!\Gamma(2-2s)}{\Gamma(4-2s+2n)} \sum_{m=0}^{\infty} \frac{(1-s)_m(n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!} I_\lambda(2m+2-2s, 2n+2).
\end{aligned}$$

By Lemma 4.11, we have

$$I_\lambda(2m+2-2s, 2n+2) = (2m+2-2s)_{2n+2} \sum_{r=0}^{2n+1} \frac{(-1)^r}{r!(2n+1-r)!} \frac{\lambda^{2m+2-2s+r}}{2m+2-2s+r}.$$

Thus,

$$\begin{aligned}
&\sum_{m=0}^{\infty} \frac{(1-s)_m(n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!} I_\lambda(2m+2-2s, 2n+2)\\
=&\sum_{r=0}^{2n+1} \frac{(-1)^r}{r!(2n+1-r)!} \sum_{m=0}^{\infty} \frac{(1-s)_m(n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!}\\
&\times \frac{(2m+2-2s)(2m+3-2s)\cdots(2m+2n+3-2s)}{2m+2-2s+r} \lambda^{2m+2-2s+r}.
\end{aligned}$$

If $n=0$, then $r=0,1$.

(1) $r=0$,

$$\sum_{m=0}^{\infty} \frac{(1-s)_m(\frac{1}{2})_m}{(\frac{5}{2}-s)_m m!} (2m+3-2s)\lambda^{2m+2-2s} = (3-2s)\lambda^{2-2s} F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2).$$

(2) $r=1$,

$$\sum_{m=0}^{\infty} \frac{(1-s)_m(\frac{1}{2})_m}{(\frac{5}{2}-s)_m m!} (2m+2-2s)\lambda^{2m+3-2s} = 2(1-s)\lambda^{3-2s} F(2-s, \frac{1}{2}; \frac{5}{2}-s; \lambda^2).$$

On the other hand, when $n=0$, (4.28) is equal to

$$\frac{1}{(3-2s)(2-2s)} F(1-s, \frac{1}{2}; \frac{5}{2}-s; 1).$$

Thus,

$$\begin{aligned}&\int_\lambda^1 (1-l)^{2n+1}l^{1-2s}F(n-s+2,n+\frac{1}{2};\frac{3}{2}-s;l^2)dl_{n=0}\\ =&\frac{1}{(3-2s)(2-2s)}F(1-s,\frac{1}{2};\frac{5}{2}-s;1)-\frac{\lambda^{2-2s}}{2-2s}F(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2)\\ &+\frac{\lambda^{3-2s}}{3-2s}F(2-s,\frac{1}{2};\frac{5}{2}-s;\lambda^2).\end{aligned}$$

If $n=1$, then $r=0,1,2,3$.
(1) $r=0$,

$$\begin{aligned}&\sum_{m=0}^\infty \frac{(1-s)_m(\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!}(2m+3-2s)(2m+4-2s)(2m+5-2s)\lambda^{2m+2-2s}\\ =&8(1-s)(\frac{3}{2}-s)(\frac{5}{2}-s)\lambda^{2-2s}F(2-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2)\\ &+8(\frac{3}{2}-s)(\frac{5}{2}-s)\lambda^{2-2s}F(1-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2).\end{aligned}$$

(2) $r=1$,

$$\begin{aligned}&\sum_{m=0}^\infty \frac{(1-s)_m(\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!}(2m+2-2s)(2m+4-2s)(2m+5-2s)\lambda^{2m+3-2s}\\ =&8(1-s)(2-s)(\frac{5}{2}-s)\lambda^{3-2s}F(3-s,\frac{3}{2};\frac{5}{2}-s;\lambda^2).\end{aligned}$$

(3) $r=2$,

$$\begin{aligned}&\sum_{m=0}^\infty \frac{(1-s)_m(\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!}(2m+2-2s)(2m+3-2s)(2m+5-2s)\lambda^{2m+4-2s}\\ =&8(1-s)(\frac{3}{2}-s)(\frac{5}{2}-s)\lambda^{4-2s}F(2-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2).\end{aligned}$$

(3) $r=3$,

$$\begin{aligned}&\sum_{m=0}^\infty \frac{(1-s)_m(\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!}(2m+2-2s)(2m+3-2s)(2m+4-2s)\lambda^{2m+5-2s}\\ =&8(1-s)(2-s)(\frac{5}{2}-s)\lambda^{5-2s}F(3-s,\frac{3}{2};\frac{5}{2}-s;\lambda^2)\\ &-8(1-s)(2-s)\lambda^{5-2s}F(3-s,\frac{3}{2};\frac{7}{2}-s;\lambda^2).\end{aligned}$$

On the other hand, when $n = 1$, (4.28) is equal to

$$\frac{6\Gamma(2-2s)}{\Gamma(6-2s)}F(1-s,\frac{3}{2};\frac{7}{2}-s;1).$$

Thus,

$$\begin{aligned}
&\int_\lambda^1 (1-l)^{2n+1}l^{1-2s}F(n-s+2,n+\frac{1}{2};\frac{3}{2}-s;l^2)_{n=1}\\
=&\frac{6\Gamma(2-2s)}{\Gamma(6-2s)}F(1-s,\frac{3}{2};\frac{7}{2}-s;1)-\frac{1}{2}\frac{1}{(2-s)}\lambda^{2-2s}(1+3\lambda^2)F(2-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2)\\
&+\frac{1}{2}\frac{1}{(\frac{3}{2}-s)}\lambda^{3-2s}(3+\lambda^2)F(3-s,\frac{3}{2};\frac{5}{2}-s;\lambda^2)-\frac{1}{2}\frac{1}{(1-s)(2-s)}\lambda^{2-2s}\\
&\times F(1-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2)-\frac{1}{2}\frac{1}{(\frac{3}{2}-s)(\frac{5}{2}-s)}\lambda^{5-2s}F(3-s,\frac{3}{2};\frac{7}{2}-s;\lambda^2).
\end{aligned}$$

Now, we need the following relation of Gauss between contiguous functions (see [Er], p.103, (33)):

$$(c-a-b)F+a(1-z)F(a+1)-(c-b)F(b-1)=0.$$

We have

$$F(1-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2)=(1-s)(1-\lambda^2)F(2-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2)+sF(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2).$$

$$F(3-s,\frac{3}{2};\frac{7}{2}-s;\lambda^2)=(3-s)(1-\lambda^2)F(4-s,\frac{3}{2};\frac{7}{2}-s;\lambda^2)-(2-s)F(3-s,\frac{1}{2};\frac{7}{2}-s;\lambda^2).$$

By [Er], p.104, (46),

$$F(a,b;c;1)=\frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)},\quad c\neq 0,-1,-2,\cdots,\quad \mathrm{Re}(c)>\mathrm{Re}(a+b),$$

we have

$$F(1-s,\frac{1}{2};\frac{5}{2}-s;1)=2\frac{\Gamma(\frac{5}{2}-s)}{\sqrt{\pi}\Gamma(2-s)},\quad F(1-s,\frac{3}{2};\frac{7}{2}-s;1)=\frac{4}{3}\frac{\Gamma(\frac{7}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}.$$

Therefore,

$$\begin{aligned}
&\int_\lambda^1 (1-l)^{2n+1}l^{1-2s}F(n-s+2,n+\frac{1}{2};\frac{3}{2}-s;l^2)dl_{n=0}\\
=&\frac{1}{2-2s}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}-\frac{\lambda^{2-2s}}{2-2s}F(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2)+\frac{\lambda^{3-2s}}{3-2s}F(2-s,\frac{1}{2};\frac{5}{2}-s;\lambda^2),
\end{aligned}$$

and

$$\begin{aligned}&\int_\lambda^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2)\, dl_{n=1}\\
=&\frac{1}{2-2s}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)} - \frac{1}{2-s}(1+\lambda^2)\lambda^{2-2s}F(2-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2)\\
&+\frac{1}{2}\frac{1}{(\frac{3}{2}-s)}(3+\lambda^2)\lambda^{3-2s}F(3-s,\frac{3}{2};\frac{5}{2}-s;\lambda^2)\\
&-\frac{1}{2}\frac{s}{(1-s)(2-s)}\lambda^{2-2s}F(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2)\\
&-\frac{1}{2}\frac{(3-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}(1-\lambda^2)\lambda^{5-2s}F(4-s,\frac{3}{2};\frac{7}{2}-s;\lambda^2)\\
&+\frac{1}{2}\frac{(2-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}\lambda^{5-2s}F(3-s,\frac{1}{2};\frac{7}{2}-s;\lambda^2).\end{aligned}$$

By [Er], p.111, (5),

$$F(a,b;2b;\frac{4z}{(1+z)^2}) = (1+z)^{2a}F(a,a+\frac{1}{2}-b;b+\frac{1}{2};z^2),$$

we have

$$\begin{aligned}F(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2) &= (1-\lambda)^{-2(1-s)}F(1-s,1-s;2-2s;-u^{-1}),\\
F(2-s,\frac{1}{2};\frac{5}{2}-s;\lambda^2) &= (1-\lambda)^{-2(2-s)}F(2-s,2-s;4-2s;-u^{-1}),\\
F(2-s,\frac{3}{2};\frac{3}{2}-s;\lambda^2) &= (1-\lambda)^{-2(2-s)}F(2-s,1-s;2-2s;-u^{-1}),\\
F(3-s,\frac{3}{2};\frac{5}{2}-s;\lambda^2) &= (1-\lambda)^{-2(3-s)}F(3-s,2-s;4-2s;-u^{-1}),\\
F(1-s,\frac{1}{2};\frac{3}{2}-s;\lambda^2) &= (1-\lambda)^{-2(1-s)}F(1-s,1-s;2-2s;-u^{-1}),\\
F(4-s,\frac{3}{2};\frac{7}{2}-s;\lambda^2) &= (1-\lambda)^{-2(4-s)}F(4-s,3-s;6-2s;-u^{-1}),\\
F(3-s,\frac{1}{2};\frac{7}{2}-s;\lambda^2) &= (1-\lambda)^{-2(3-s)}F(3-s,3-s;6-2s;-u^{-1}).\end{aligned}$$

Therefore,

$$\begin{aligned}&\int_\lambda^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2)\, dl_{n=0}\\
=&\frac{1}{2-2s}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} - \frac{1}{2-2s}\lambda^{2-2s}(1-\lambda)^{-2(1-s)}F(1-s,1-s;2-2s;-u^{-1})\\
&+\frac{1}{3-2s}\lambda^{3-2s}(1-\lambda)^{-2(2-s)}F(2-s,2-s;4-2s;-u^{-1}).\end{aligned}$$

$$\begin{aligned}
&\int_\lambda^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2)dl_{n=1} \\
=&\frac{1}{2-2s}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)} - \frac{1}{2-s}(1+\lambda^2)\lambda^{2-2s}(1-\lambda)^{-2(2-s)}F(2-s,1-s;2-2s;-u^{-1}) \\
&+\frac{1}{2}\frac{1}{(\frac{3}{2}-s)}(3+\lambda^2)\lambda^{3-2s}(1-\lambda)^{-2(3-s)}F(3-s,2-s;4-2s;-u^{-1}) \\
&-\frac{1}{2}\frac{s}{(1-s)(2-s)}\lambda^{2-2s}(1-\lambda)^{-2(1-s)}F(1-s,1-s;2-2s;-u^{-1}) \\
&-\frac{1}{2}\frac{(3-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}(1-\lambda^2)\lambda^{5-2s}(1-\lambda)^{-2(4-s)}F(4-s,3-s;6-2s;-u^{-1}) \\
&+\frac{1}{2}\frac{(2-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}\lambda^{5-2s}(1-\lambda)^{-2(3-s)}F(3-s,3-s;6-2s;-u^{-1}).
\end{aligned}$$

Since $u=\frac{(\lambda-1)^2}{4\lambda}$, we have

$$\begin{aligned}
&\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}\lambda^s(1-\lambda)^{-2}\int_\lambda^1 (1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=0} \\
=&\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}[-(4u)^{-(s+1)}F(s,s;2s;-u^{-1})+\frac{s-\frac{1}{2}}{s-1}(4u)^{-s} \\
&\times F(s-1,s-1;2s-2;-u^{-1})-\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}\lambda^s(1-\lambda)^{-2}],
\end{aligned}$$

and

$$\begin{aligned}
&\frac{2\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}\lambda^s(1-\lambda)^{-2}\int_\lambda^1 (1-l)^{2n+1}l^{1-2s}F(n-s+2,n+\frac{1}{2};\frac{3}{2}-s;l^2)dl_{n=0} \\
=&\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}[\frac{1}{1-s}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}\lambda^s(1-\lambda)^{-2}-\frac{1}{1-s}(4u)^{-(2-s)} \\
&\times F(1-s,1-s;2-2s;-u^{-1})+\frac{1}{\frac{3}{2}-s}(4u)^{-(3-s)}F(2-s,2-s;4-2s;-u^{-1})].
\end{aligned}$$

Note that

$$\begin{aligned}
&\lambda^s(1-\lambda)^{-2}\left[\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}\frac{(-1)}{\sqrt{\pi}}\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\Gamma(s)}+\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}\frac{1}{\sqrt{\pi}}\frac{1}{1-s}\frac{\Gamma(\frac{3}{2}-s)}{\Gamma(2-s)}\right] \\
=&\frac{\lambda^s(1-\lambda)^{-2}}{\pi(1-s)\Gamma(s)\Gamma(2-s)}[\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2})+\Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)] \\
=&0.
\end{aligned}$$

Now, we need the following formulas:

$$\gamma F-\beta zF(\beta+1,\gamma+1)-\gamma F(\alpha-1)=0.$$

$$(\gamma-1)F(\gamma-1)-\alpha F(\alpha+1)-(\gamma-\alpha-1)F=0.$$

We have

$$(2s-1)F(s,s-1;2s-1;z)-(s-1)zF(s,s;2s;z)-(2s-1)F(s-1,s-1;2s-1;z)=0,$$

and

$$(2s-2)F(s-1,s-1;2s-2;z)-(s-1)F(s,s-1;2s-1;z)-(s-1)F(s-1,s-1;2s-1;z)=0.$$

Thus,

$$\frac{4s-2}{s-1}F(s,s-1;2s-1;z)=zF(s,s;2s;z)+\frac{4s-2}{s-1}F(s-1,s-1;2s-2;z).$$

Hence, we have

$$\begin{aligned}&\frac{s-\frac{1}{2}}{s-1}F(s,s-1;2s-1;-u^{-1})\\ =&-(4u)^{-1}F(s,s;2s;-u^{-1})+\frac{s-\frac{1}{2}}{s-1}F(s-1,s-1;2s-2;-u^{-1}).\end{aligned}$$

Similarly,

$$\begin{aligned}&\frac{\frac{3}{2}-s}{1-s}F(2-s,1-s;3-2s;-u^{-1})\\ =&-(4u)^{-1}F(2-s,2-s;4-2s;-u^{-1})+\frac{\frac{3}{2}-s}{1-s}F(1-s,1-s;2-2s;-u^{-1}).\end{aligned}$$

Therefore,

$$\begin{aligned}\frac{1}{4\pi^2}\phi(\frac{1}{2}\rho,\frac{1}{2}\rho';0,s)=&\frac{1}{1-s}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}(4u)^{-s}F(s,s-1;2s-1;-u^{-1})\\ &+\frac{1}{s-1}\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}(4u)^{s-2}F(2-s,1-s;3-2s;-u^{-1}).\end{aligned}$$

For $n=1$, note that

$$\frac{3!\Gamma(\frac{1}{2}-s)}{\Gamma(3-s)\Gamma(\frac{3}{2})}=\frac{12\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)},$$

we have

$$\begin{aligned}
&\frac{12\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\lambda^s(1-\lambda)^{-2}\int_\lambda^1(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=1}\\
=&-4\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}\lambda^s(1-\lambda)^{-2}\\
&-4\frac{\Gamma(\frac{5}{2}-s)}{s(s-1)\sqrt{\pi}\Gamma(3-s)}\lambda(4u)^{-(s-1)}F(s-2,s-2;2s-4;-u^{-1})\\
&-4\frac{\Gamma(\frac{5}{2}-s)}{s(s-1)\sqrt{\pi}\Gamma(3-s)}(1-\lambda^2)(4u)^{-s}F(s-1,s-2;2s-4;-u^{-1})\\
&-4\frac{\Gamma(\frac{3}{2}-s)}{s\sqrt{\pi}\Gamma(3-s)}(\lambda+\frac{3}{\lambda})(4u)^{-(s+1)}F(s,s-1;2s-2;-u^{-1})\\
&-4\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}(4u)^{-(s+1)}F(s,s;2s;-u^{-1})\\
&-8\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}(\lambda+\frac{1}{\lambda})(4u)^{-(s+2)}F(s+1,s;2s;-u^{-1}).
\end{aligned}$$

Similarly,

$$\frac{3!\Gamma(s-\frac{1}{2})}{\Gamma(s+1)\Gamma(\frac{5}{2})}=\frac{8\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)},$$

and

$$\begin{aligned}
&\frac{8\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}\lambda^s(1-\lambda)^{-2}\int_\lambda^1(1-l)^{2n+1}l^{1-2s}F(n-s+2,n+\frac{1}{2};\frac{3}{2}-s;l^2)dl_{n=1}\\
=&\frac{8\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}\times[\frac{1}{2}\frac{1}{(1-s)}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\lambda^s(1-\lambda)^{-2}\\
&-\frac{1}{2-s}(\lambda+\frac{1}{\lambda})(4u)^{-(3-s)}F(2-s,1-s;2-2s;-u^{-1})\\
&+\frac{1}{2}\frac{1}{(\frac{3}{2}-s)}(\lambda+\frac{3}{\lambda})(4u)^{-(4-s)}F(3-s,2-s;4-2s;-u^{-1})\\
&-\frac{1}{2}\frac{s}{(1-s)(2-s)}(4u)^{-(2-s)}F(1-s,1-s;2-2s;-u^{-1})\\
&-\frac{1}{2}\frac{(3-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}(1-\lambda^2)(4u)^{-(5-s)}F(4-s,3-s;6-2s;-u^{-1})\\
&+\frac{1}{2}\frac{(2-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}\lambda(4u)^{-(4-s)}F(3-s,3-s;6-2s;-u^{-1})].
\end{aligned}$$

Note that

$$\begin{aligned}&\left[\frac{-4\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}+\frac{-4\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}\frac{1}{s-1}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\right]\lambda^s(1-\lambda)^{-2}\\ =&\frac{-4\lambda^s(1-\lambda)^{-2}}{\pi(s-1)\Gamma(s+1)\Gamma(3-s)}\left[\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2})+\Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)\right]\\ =&0.\end{aligned}$$

For $k=1$, if $n=0$, $\frac{(-1)_n(1)_n}{2^{2n}(n!)^2}=1$. If $n=1$, $\frac{(-1)_n(1)_n}{2^{2n}(n!)^2}=-\frac{1}{4}$. Let us add the terms of $n=0$ and $n=1$, we have

$$\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}(4u)^{-s}S_I+\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}(4u)^{s-3}S_{II},$$

where

$$\begin{aligned}S_I=&\frac{\frac{1}{2}-s}{1-s}F(s,s-1;2s-1;-u^{-1})+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}4u\lambda F(s-2,s-2;2s-4;-u^{-1})\\ &+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}(1-\lambda^2)F(s-1,s-2;2s-4;-u^{-1})\\ &+\frac{\frac{1}{2}-s}{s(2-s)}(\lambda+\frac{3}{\lambda})\frac{1}{4u}F(s,s-1;2s-2;-u^{-1})\\ &+\frac{1}{2-s}\frac{1}{4u}F(s,s;2s;-u^{-1})+\frac{1}{2-s}2(\lambda+\frac{1}{\lambda})\frac{1}{(4u)^2}F(s+1,s;2s;-u^{-1}),\end{aligned}$$

and

$$\begin{aligned}S_{II}=&\frac{1}{s-1}4uF(2-s,1-s;3-2s;-u^{-1})\\ &+\frac{2}{s(2-s)}(\lambda+\frac{1}{\lambda})F(2-s,1-s;2-2s;-u^{-1})\\ &-\frac{1}{s(\frac{3}{2}-s)}(\lambda+\frac{3}{\lambda})(4u)^{-1}F(3-s,2-s;4-2s;-u^{-1})\\ &+\frac{1}{(1-s)(2-s)}4uF(1-s,1-s;2-2s;-u^{-1})\\ &+\frac{(3-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)}(1-\lambda^2)(4u)^{-2}F(4-s,3-s;6-2s;-u^{-1})\\ &-\frac{(2-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)}\lambda(4u)^{-1}F(3-s,3-s;6-2s;-u^{-1}).\end{aligned}$$

Since $u = \frac{(\lambda-1)^2}{4\lambda}$, we have

$$\lambda = \frac{1}{(\sqrt{u+1}+\sqrt{u})^2}.$$

Hence,

$$\lambda + \frac{1}{\lambda} = 4u+2, \quad \lambda + \frac{3}{\lambda} = 4(2u+1+\sqrt{u+1}\sqrt{u}),$$
$$1-\lambda^2 = \lambda(\frac{1}{\lambda}-\lambda) = \frac{4\sqrt{u}\sqrt{u+1}}{(\sqrt{u+1}+\sqrt{u})^2}.$$

Thus, we have

$$\begin{aligned}
S_I =& \frac{s-\frac{1}{2}}{s-1}F(s,s-1;2s-1;-u^{-1}) \\
&+ \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{4}{(\sqrt{1+\frac{1}{u}}+1)^2}F(s-2,s-2;2s-4;-u^{-1}) \\
&+ \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{4\sqrt{1+\frac{1}{u}}}{(\sqrt{1+\frac{1}{u}}+1)^2}F(s-1,s-2;2s-4;-u^{-1}) \\
&+ \frac{(\frac{1}{2}-s)}{s(2-s)}(2+\frac{1}{u}+\sqrt{1+\frac{1}{u}})F(s,s-1;2s-2;-u^{-1}) \\
&+ \frac{1}{2-s}\frac{1}{4u}F(s,s;2s;-u^{-1}) + \frac{1}{2-s}(\frac{1}{2u}+\frac{1}{4u^2})F(s+1,s;2s;-u^{-1}).
\end{aligned}$$

$$\begin{aligned}
S_{II} =& \frac{1}{s-1}4uF(2-s,1-s;3-2s;-u^{-1}) \\
&+ \frac{1}{s(2-s)}4(2u+1)F(2-s,1-s;2-2s;-u^{-1}) \\
&- \frac{1}{s(\frac{3}{2}-s)}(2+\frac{1}{u}+\sqrt{1+\frac{1}{u}})F(3-s,2-s;4-2s;-u^{-1}) \\
&+ \frac{1}{(1-s)(2-s)}4uF(1-s,1-s;2-2s;-u^{-1}) \\
&+ \frac{(3-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)}\frac{\sqrt{1+\frac{1}{u}}}{(1+\sqrt{1+\frac{1}{u}})^2}\frac{1}{4u^2}F(4-s,3-s;6-2s;-u^{-1}) \\
&- \frac{(2-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)}\frac{1}{(1+\sqrt{1+\frac{1}{u}})^2}\frac{1}{4u^2}F(3-s,3-s;6-2s;-u^{-1}).
\end{aligned}$$

Set $z = -u^{-1}$ and $x = \sqrt{1-z}$, then

$$\frac{u}{u+1} = \frac{1}{x^2}, \quad \frac{u^2}{u+1} = \frac{1}{x^2(x^2-1)}.$$

$$\begin{aligned} S_I = & \frac{s-\frac{1}{2}}{s-1} F(s, s-1; 2s-1; 1-x^2) \\ & + \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)} \frac{4}{(x+1)^2} F(s-2, s-2; 2s-4; 1-x^2) \\ & + \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)} \frac{4x}{(x+1)^2} F(s-1, s-2; 2s-4; 1-x^2) \\ & + \frac{(\frac{1}{2}-s)}{s(2-s)} (x^2+x+1) F(s, s-1; 2s-2; 1-x^2) \\ & + \frac{1}{2-s} \frac{1}{4} (x^2-1) F(s, s; 2s; 1-x^2) + \frac{1}{2-s} \frac{1}{4} (x^4-1) F(s+1, s; 2s; 1-x^2). \end{aligned}$$

$$\begin{aligned} S_{II} = & \frac{1}{s-1} \frac{4}{x^2-1} F(2-s, 1-s; 3-2s; 1-x^2) \\ & + \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2-1} F(2-s, 1-s; 2-2s; 1-x^2) \\ & - \frac{1}{s(\frac{3}{2}-s)} (x^2+x+1) F(3-s, 2-s; 4-2s; 1-x^2) \\ & + \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} F(1-s, 1-s; 2-2s; 1-x^2) \\ & + \frac{(3-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)} \frac{1}{4} x(x-1)^2 F(4-s, 3-s; 6-2s; 1-x^2) \\ & - \frac{(2-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)} \frac{1}{4} (x-1)^2 F(3-s, 3-s; 6-2s; 1-x^2). \end{aligned}$$

At first, we calculate S_I. It is known that (see [Er], p.103, (29))

$$(c-b)F(b-1) + (2b-c-bz+az)F + b(z-1)F(b+1) = 0.$$

Set $a = s$, $b = s$, $c = 2s$ and $z = 1-x^2$, then

$$F(s, s+1; 2s; 1-x^2) = \frac{1}{x^2} F(s, s-1; 2s; 1-x^2),$$

i.e.,

$$F(s+1, s; 2s; 1-x^2) = \frac{1}{x^2} F(s, s-1; 2s; 1-x^2).$$

Set $a = s-1$, $b = s-1$, $c = 2s-2$ and $z = 1-x^2$, then

$$F(s-1,s;2s-2;1-x^2) = \frac{1}{x^2}F(s-1,s-2;2s-2;1-x^2),$$

i.e.,

$$F(s,s-1;2s-2;1-x^2) = \frac{1}{x^2}F(s-1,s-2;2s-2;1-x^2).$$

In the formula (see [Er], p.103, (34))

$$c[a-(c-b)z]F - ac(1-z)F(a+1) + (c-a)(c-b)zF(c+1) = 0,$$

set $a = s-2$, $b = s-2$, $c = 2s-4$ and $z = 1-x^2$, then

$$\begin{aligned}&F(s-2,s-2;2s-4;1-x^2)\\ =&F(s-1,s-2;2s-4;1-x^2) + \frac{x^2-1}{2x^2}F(s-2,s-2;2s-3;1-x^2).\end{aligned}$$

Now, we have

$$\begin{aligned}S_I =&\frac{s-\frac{1}{2}}{s-1}F(s,s-1;2s-1;1-x^2)\\ &+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{4}{x+1}F(s-1,s-2;2s-4;1-x^2)\\ &+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{2(x-1)}{x^2(x+1)}F(s-2,s-2;2s-3;1-x^2)\\ &+\frac{(\frac{1}{2}-s)}{s(2-s)}\frac{x^2+x+1}{x^2}F(s-1,s-2;2s-2;1-x^2)\\ &+\frac{1}{2-s}\frac{1}{4}(x^2-1)F(s,s;2s;1-x^2)\\ &+\frac{1}{2-s}\frac{1}{4}\frac{x^4-1}{x^2}F(s,s-1;2s;1-x^2).\end{aligned}$$

It is known that

$$\gamma F - \beta z F(\beta+1,\gamma+1) - \gamma F(\alpha-1) = 0.$$

Set $\alpha = s$, $\beta = s-1$, $\gamma = 2s-1$ and $z = 1-x^2$, then

$$F(s,s;2s;1-x^2) = \frac{1}{1-x^2}\frac{2s-1}{s-1}[F(s,s-1;2s-1;1-x^2) - F(s-1,s-1;2s-1;1-x^2)].$$

Set $\alpha = s-1$, $\beta = s-1$, $\gamma = 2s-1$ and $z = 1-x^2$, then

$$\begin{aligned}
&F(s,s-1;2s;1-x^2) = F(s-1,s;2s;1-x^2)\\
=&\frac{1}{1-x^2}\frac{2s-1}{s-1}[F(s-1,s-1;2s-1;1-x^2) - F(s-2,s-1;2s-1;1-x^2)].
\end{aligned}$$

Thus, we have

$$\begin{aligned}
S_I =&\frac{(\frac{1}{2}-s)(s-\frac{3}{2})}{(2-s)(s-1)}F(s,s-1;2s-1;1-x^2)\\
&+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{4}{x+1}F(s-1,s-2;2s-4;1-x^2)\\
&+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{2(x-1)}{x^2(x+1)}F(s-2,s-2;2s-3;1-x^2)\\
&+\frac{(\frac{1}{2}-s)}{s(2-s)}\frac{x^2+x+1}{x^2}F(s-1,s-2;2s-2;1-x^2)\\
&+\frac{(\frac{1}{2}-s)}{(2-s)(s-1)}\frac{1}{2x^2}F(s-1,s-1;2s-1;1-x^2)\\
&-\frac{(\frac{1}{2}-s)}{(2-s)(s-1)}\frac{x^2+1}{2x^2}F(s-1,s-2;2s-1;1-x^2).
\end{aligned}$$

It is known that (see [Er], p.103, (37))

$$(b-a)(1-z)F - (c-a)F(a-1) + (c-b)F(b-1) = 0.$$

Set $a = s$, $b = s-1$, $c = 2s-1$ and $z = 1-x^2$, then

$$F(s,s-1;2s-1;1-x^2) = \frac{1-s}{x^2}F(s-1,s-1;2s-1;1-x^2) + \frac{s}{x^2}F(s,s-2;2s-1;1-x^2).$$

By [Er], p.103, (30),

$$c(c-1)(z-1)F(c-1) + c[c-1-(2c-a-b-1)z]F + (c-a)(c-b)zF(c+1) = 0,$$

set $a = s-1$, $b = s-2$, $c = 2s-3$ and $z = 1-x^2$, then

$$\begin{aligned}
&F(s-1,s-2;2s-4;1-x^2)\\
=&F(s-1,s-2;2s-3;1-x^2) + \frac{(s-1)}{4(s-\frac{3}{2})}\frac{1-x^2}{x^2}F(s-1,s-2;2s-2;1-x^2).
\end{aligned}$$

Hence,

$$\begin{aligned}S_I =&\frac{s(\frac{1}{2}-s)(\frac{3}{2}-s)}{(2-s)(1-s)}\frac{1}{x^2}F(s,s-2;2s-1;1-x^2)\\&+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{4}{x+1}F(s-1,s-2;2s-3;1-x^2)\\&+\frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)}\frac{2(x-1)}{x^2(x+1)}F(s-2,s-2;2s-3;1-x^2)\\&+\frac{(\frac{1}{2}-s)}{s(2-s)}\frac{x+2}{x}F(s-1,s-2;2s-2;1-x^2)\\&-\frac{(\frac{1}{2}-s)^2}{s-1}\frac{1}{x^2}F(s-1,s-1;2s-1;1-x^2)\\&-\frac{(\frac{1}{2}-s)}{(2-s)(s-1)}\frac{x^2+1}{2x^2}F(s-1,s-2;2s-1;1-x^2).\end{aligned}$$

It is known that (see [Er], p.103, (35))

$$(c-a-1)F+aF(a+1)-(c-1)F(c-1)=0.$$

Set $a=s-2$, $b=s-1$, $c=2s-1$ and $z=1-x^2$, then

$$\begin{aligned}&F(s-1,s-1;2s-1;1-x^2)\\=&\frac{2(s-1)}{s-2}F(s-2,s-1;2s-2;1-x^2)-\frac{s}{s-2}F(s-2,s-1;2s-1;1-x^2).\end{aligned}$$

Set $a=s-2$, $b=s-2$, $c=2s-2$ and $z=1-x^2$, then

$$\begin{aligned}&F(s-2,s-2;2s-3;1-x^2)\\=&\frac{(s-1)}{2(s-\frac{3}{2})}F(s-2,s-2;2s-2;1-x^2)+\frac{(s-2)}{2(s-\frac{3}{2})}F(s-1,s-2;2s-2;1-x^2).\end{aligned}$$

Set $a=s-1$, $b=s-2$, $c=2s-1$ and $z=1-x^2$, then

$$F(s,s-2;2s-1;1-x^2)=2F(s-1,s-2;2s-2;1-x^2)-F(s-1,s-2;2s-1;1-x^2).$$

Hence,

$$\begin{aligned}
S_I = &\left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2}+\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{x^2+1}{2x^2}\right]F(s-1,s-2;2s-1;1-x^2)\\
&+\left[\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2}+\frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{x^2(x+1)}-\frac{(\frac{1}{2}-s)}{s(s-2)}\frac{x+2}{x}\right]\\
&\times F(s-1,s-2;2s-2;1-x^2)\\
&+\frac{(\frac{1}{2}-s)(\frac{3}{2}-s)}{s(s-1)(2-s)}\frac{4}{x+1}F(s-1,s-2;2s-3;1-x^2)\\
&+\frac{(\frac{1}{2}-s)}{s(s-2)}\frac{(x-1)}{x^2(x+1)}F(s-2,s-2;2s-2;1-x^2).
\end{aligned}$$

We know that (see [Er], p.103, (35))

$$(c-a-1)F+aF(a+1)-(c-1)F(c-1)=0.$$

Set $a=s-2$, $b=s-2$, $c=2s-1$ and $z=1-x^2$, then

$$\begin{aligned}
&F(s-2,s-2;2s-2;1-x^2)\\
=&\frac{s}{2(s-1)}F(s-2,s-2;2s-1;1-x^2)+\frac{s-2}{2(s-1)}F(s-1,s-2;2s-1;1-x^2).
\end{aligned}$$

For $F(s-1,s-2;2s-3;1-x^2)=F(s-2,s-1;2s-3;1-x^2)$, set $a=s-2$, $b=s-1$, $c=2s-2$ and $z=1-x^2$, then

$$\begin{aligned}
&F(s-1,s-2;2s-3;1-x^2)\\
=&\frac{s-1}{2(s-\frac{3}{2})}F(s-1,s-2;2s-2;1-x^2)+\frac{s-2}{2(s-\frac{3}{2})}F(s-1,s-1;2s-2;1-x^2).
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
S_I = &\left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2}+\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{x^2+1}{2x^2}+\frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{2x^2(x+1)}\right]\\
&\times F(s-1,s-2;2s-1;1-x^2)\\
&+\left[\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2}+\frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{x^2(x+1)}-\frac{(\frac{1}{2}-s)}{s(s-2)}\frac{(x+2)}{x}+\frac{(\frac{1}{2}-s)}{s(s-2)}\frac{2}{x+1}\right]\\
&\times F(s-1,s-2;2s-2;1-x^2)\\
&+\frac{(\frac{1}{2}-s)}{s(s-1)}\frac{2}{x+1}F(s-1,s-1;2s-2;1-x^2)\\
&+\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x-1)}{2x^2(x+1)}F(s-2,s-2;2s-1;1-x^2).
\end{aligned}$$

Since (see [Er], p.103, (38))

$$c(1-z)F - cF(a-1) + (c-b)zF(c+1) = 0.$$

Set $a = s-1$, $b = s-1$, $c = 2s-2$ and $z = 1-x^2$, then

$$\begin{aligned}&F(s-1, s-1; 2s-2; 1-x^2)\\ =&\frac{1}{x^2}F(s-1, s-2; 2s-2; 1-x^2) + \frac{x^2-1}{2x^2}F(s-1, s-1; 2s-1; 1-x^2).\end{aligned}$$

Thus,

$$\begin{aligned}S_I = &\left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x^2+1)}{2x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{2x^2(x+1)}\right]\\ &\times F(s-1, s-2; 2s-1; 1-x^2)\\ &+ \frac{(\frac{1}{2}-s)}{s(s-2)}\frac{-(x-1)(x^2+2x+2)}{x^2(x+1)}F(s-1, s-2; 2s-2; 1-x^2)\\ &+ \frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{x^2}F(s-1, s-1; 2s-1; 1-x^2)\\ &+ \frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x-1)}{2x^2(x+1)}F(s-2, s-2; 2s-1; 1-x^2).\end{aligned}$$

It is known that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = s-2$, $b = s-1$, $c = 2s-1$ and $z = 1-x^2$, then

$$\begin{aligned}&F(s-1, s-2; 2s-2; 1-x^2)\\ =&\frac{s}{2(s-1)}F(s-1, s-2; 2s-1; 1-x^2) + \frac{(s-2)}{2(s-1)}F(s-1, s-1; 2s-1; 1-x^2).\end{aligned}$$

Hence,

$$\begin{aligned}S_I =&[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x^2+1)}{2x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{2x^2(x+1)}\\ &+ \frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{-(x-1)(x^2+2x+2)}{2x^2(x+1)}]F(s-1, s-2; 2s-1; 1-x^2)\\ &+ \frac{(\frac{1}{2}-s)}{s(s-1)}\frac{-(x-1)}{2(x+1)}F(s-1, s-1; 2s-1; 1-x^2)\\ &+ \frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x-1)}{2x^2(x+1)}F(s-2, s-2; 2s-1; 1-x^2).\end{aligned} \tag{4.30}$$

In the following formula (see [Er], p.103, (36))

$$(c-a-b)F-(c-a)F(a-1)+b(1-z)F(b+1)=0,$$

set $a=s-1$, $b=s-2$, $c=2s-1$ and $z=1-x^2$, then

$$\begin{aligned}&2F(s-1,s-2;2s-1;1-x^2)\\=&-(s-2)x^2F(s-1,s-1,2s-1;1-x^2)+sF(s-2,s-2;2s-1;1-x^2).\end{aligned}$$

Now, the sum of the latter two terms in (4.30) is equal to

$$\begin{aligned}&\frac{(\frac{1}{2}-s)}{s(s-1)(s-2)}\frac{(x-1)}{2x^2(x+1)}[-(s-2)x^2F(s-1,s-1;2s-1;1-x^2)\\&+sF(s-2,s-2;2s-1;1-x^2)]\\=&\frac{(\frac{1}{2}-s)}{s(s-1)(s-2)}\frac{(x-1)}{x^2(x+1)}F(s-1,s-2;2s-1;1-x^2).\end{aligned}$$

Thus,

$$\begin{aligned}S_I=&[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2}+\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x^2+1)}{2x^2}+\frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{2x^2(x+1)}\\&+\frac{(\frac{1}{2}-s)}{s(s-1)(s-2)}\frac{(x-1)}{x^2(x+1)}+\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{-(x-1)(x^2+2x+2)}{2x^2(x+1)}]\\&\times F(s-1,s-2;2s-1;1-x^2).\end{aligned}$$

Note that

$$\frac{x^2+1}{2x^2}-\frac{(x-1)(x^2+2x+2)}{2x^2(x+1)}=\frac{x+3}{2x^2(x+1)},$$

and

$$\frac{(\frac{1}{2}-s)}{s(s-1)}\frac{(x-1)}{2x^2(x+1)}+\frac{(\frac{1}{2}-s)}{s(s-1)(s-2)}\frac{(x-1)}{x^2(x+1)}=\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x-1)}{2x^2(x+1)}.$$

We have

$$\begin{aligned}S_I=&\left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{1}{x^2}+\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x+3)}{2x^2(x+1)}+\frac{(\frac{1}{2}-s)}{(s-1)(s-2)}\frac{(x-1)}{2x^2(x+1)}\right]\\&\times F(s-1,s-2;2s-1;1-x^2)\\=&\frac{\frac{1}{2}-s}{2-s}\frac{1}{x^2}F(s-1,s-2;2s-1;1-x^2)\\=&\frac{\frac{1}{2}-s}{2-s}\frac{u}{u+1}F(s-1,s-2;2s-1;-u^{-1}).\end{aligned}$$

Therefore, we have

$$\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}(4u)^{-s}S_I = \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}4^{-s}(u+1)^{-1}u^{-(s-1)}F(s-1,s-2;2s-1;-u^{-1}).$$

Now, we compute S_{II}. It is known that (see [Er], p.103, (28))

$$(c-a)F(a-1)+(2a-c-az+bz)F+a(z-1)F(a+1)=0.$$

Set $a=3-s$, $b=3-s$, $c=6-2s$ and $z=1-x^2$, then

$$F(4-s,3-s;6-2s;1-x^2)=\frac{1}{x^2}F(2-s,3-s;6-2s;1-x^2).$$

By [Er], p.103, (39),

$$[a-1-(c-b-1)z]F+(c-a)F(a-1)-(c-1)(1-z)F(c-1)=0,$$

set $a=3-s$, $b=3-s$, $c=6-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(3-s,3-s;6-2s;1-x^2)\\=&\frac{2(\frac{5}{2}-s)}{2-s}F(3-s,3-s;5-2s;1-x^2)-\frac{3-s}{2-s}\frac{1}{x^2}F(2-s,3-s;6-2s;1-x^2).\end{aligned}$$

Thus,

$$\begin{aligned}S_{II}=&\frac{1}{s-1}\frac{4}{x^2-1}F(2-s,1-s;3-2s;1-x^2)\\&+\frac{1}{s(2-s)}\frac{4(x^2+1)}{x^2-1}F(2-s,1-s;2-2s;1-x^2)\\&-\frac{1}{s(\frac{3}{2}-s)}(x^2+x+1)F(3-s,2-s;4-2s;1-x^2)\\&+\frac{1}{(1-s)(2-s)}\frac{4}{x^2-1}F(1-s,1-s;2-2s;1-x^2)\\&+\frac{(3-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)}\frac{(x-1)^2(x+1)}{4x^2}F(3-s,2-s;6-2s;1-x^2)\\&-\frac{1}{s(\frac{3}{2}-s)}\frac{1}{2}(x-1)^2F(3-s,3-s;5-2s;1-x^2).\end{aligned}$$

We know that (see [Er], p.103, (30))

$$c(c-1)(z-1)F(c-1)+c[c-1-(2c-a-b-1)z]F+(c-a)(c-b)zF(c+1)=0.$$

Set $a = 3 - s$, $b = 2 - s$, $c = 5 - 2s$ and $z = 1 - x^2$, then

$$\begin{aligned} F(3-s, 2-s; 6-2s; 1-x^2) =& \frac{-4(\frac{5}{2}-s)}{3-s} \frac{x^2}{x^2-1} F(3-s, 2-s; 4-2s; 1-x^2) \\ &+ \frac{4(\frac{5}{2}-s)}{3-s} \frac{x^2}{x^2-1} F(3-s, 2-s; 5-2s; 1-x^2). \end{aligned}$$

It is known that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = 2 - s$, $b = 3 - s$, $c = 5 - 2s$ and $z = 1 - x^2$, then

$$F(3-s, 3-s; 5-2s; 1-x^2) = 2F(3-s, 2-s; 4-2s; 1-x^2) - F(3-s, 2-s; 5-2s; 1-x^2).$$

Hence,

$$\begin{aligned} S_{II} =& \frac{1}{s-1} \frac{4}{x^2-1} F(2-s, 1-s; 3-2s; 1-x^2) \\ &+ \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2-1} F(2-s, 1-s; 2-2s; 1-x^2) \\ &- \frac{1}{s(\frac{3}{2}-s)} (2x^2+1) F(3-s, 2-s; 4-2s; 1-x^2) \\ &+ \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} F(1-s, 1-s; 2-2s; 1-x^2) \\ &+ \frac{1}{s(\frac{3}{2}-s)} \frac{1}{2} (x^2-1) F(3-s, 2-s; 5-2s; 1-x^2). \end{aligned}$$

By [Er], p.103, (39)

$$[a-1-(c-b-1)z]F + (c-a)F(a-1) - (c-1)(1-z)F(c-1) = 0,$$

set $a = 3 - s$, $b = 2 - s$, $c = 5 - 2s$ and $z = 1 - x^2$, then

$$F(3-s, 2-s; 5-2s; 1-x^2) = -\frac{1}{x^2} F(2-s, 2-s; 5-2s; 1-x^2) + 2F(3-s, 2-s; 4-2s; 1-x^2).$$

We have

$$\begin{aligned} S_{II} =& \frac{1}{s-1} \frac{4}{x^2-1} F(2-s, 1-s; 3-2s; 1-x^2) \\ &+ \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2-1} F(2-s, 1-s; 2-2s; 1-x^2) \\ &- \frac{1}{s(\frac{3}{2}-s)} (x^2+2) F(3-s, 2-s; 4-2s; 1-x^2) \\ &+ \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} F(1-s, 1-s; 2-2s; 1-x^2) \\ &- \frac{1}{s(\frac{3}{2}-s)} \frac{x^2-1}{2x^2} F(2-s, 2-s; 5-2s; 1-x^2). \end{aligned}$$

It is known that (see [Er], p.103, (28))

$$(c-a)F(a-1)+(2a-c-az+bz)F+a(z-1)F(a+1)=0.$$

Set $a=2-s$, $b=2-s$, $c=4-2s$ and $z=1-x^2$, then

$$F(3-s,2-s;4-2s;1-x^2)=\frac{1}{x^2}F(2-s,1-s;4-2s;1-x^2).$$

By

$$\gamma F-\beta zF(\beta+1,\gamma+1)-\gamma F(\alpha-1)=0,$$

set $\alpha=2-s$, $\beta=1-s$, $\gamma=4-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(2-s,2-s;5-2s;1-x^2)\\=&\frac{2(2-s)}{1-s}\frac{1}{x^2-1}[F(1-s,1-s;4-2s;1-x^2)-F(2-s,1-s;4-2s;1-x^2)].\end{aligned}$$

So,

$$\begin{aligned}S_{II}=&\frac{1}{s-1}\frac{4}{x^2-1}F(2-s,1-s;3-2s;1-x^2)\\&+\frac{1}{s(2-s)}\frac{4(x^2+1)}{x^2-1}F(2-s,1-s;2-2s;1-x^2)\\&+\left[\frac{1}{(1-s)(\frac{3}{2}-s)}\frac{1}{x^2}-\frac{1}{s(\frac{3}{2}-s)}\right]F(2-s,1-s;4-2s;1-x^2)\\&+\frac{1}{(1-s)(2-s)}\frac{4}{x^2-1}F(1-s,1-s;2-2s;1-x^2)\\&-\frac{(2-s)}{s(1-s)(\frac{3}{2}-s)}\frac{1}{x^2}F(1-s,1-s;4-2s;1-x^2).\end{aligned}$$

We know that (see [Er], p.103, (38))

$$c(1-z)F-cF(a-1)+(c-b)zF(c+1)=0.$$

Set $a=2-s$, $b=1-s$, $c=2-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(2-s,1-s;2-2s;1-x^2)\\=&\frac{1}{x^2}F(1-s,1-s;2-2s;1-x^2)+\frac{x^2-1}{2x^2}F(2-s,1-s;3-2s;1-x^2).\end{aligned}$$

Thus,

$$\begin{aligned}S_{II} =& \left[\frac{1}{s-1}\frac{4}{x^2-1}+\frac{1}{s(2-s)}\frac{2(x^2+1)}{x^2}\right]F(2-s,1-s;3-2s;1-x^2)\\&+\left[\frac{1}{(1-s)(2-s)}\frac{4}{x^2-1}+\frac{1}{s(2-s)}\frac{4(x^2+1)}{x^2(x^2-1)}\right]F(1-s,1-s;2-2s;1-x^2)\\&+\left[\frac{1}{(1-s)(\frac{3}{2}-s)}\frac{1}{x^2}-\frac{1}{s(\frac{3}{2}-s)}\right]F(2-s,1-s;4-2s;1-x^2)\\&-\frac{(2-s)}{s(1-s)(\frac{3}{2}-s)}\frac{1}{x^2}F(1-s,1-s;4-2s;1-x^2).\end{aligned}$$

It is known that

$$\gamma F-\beta zF(\beta+1,\gamma+1)-\gamma F(\alpha-1)=0.$$

Set $\alpha=1-s$, $\beta=-s$, $\gamma=3-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(1-s,1-s;4-2s;1-x^2)\\=&\frac{2(\frac{3}{2}-s)}{s}\frac{1}{x^2-1}[F(1-s,-s;3-2s;1-x^2)-F(-s,-s;3-2s;1-x^2)].\end{aligned}$$

Set $\alpha=2-s$, $\beta=-s$, $\gamma=3-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(2-s,1-s;4-2s;1-x^2)\\=&\frac{2(\frac{3}{2}-s)}{s}\frac{1}{x^2-1}[F(2-s,-s;3-2s;1-x^2)-F(1-s,-s;3-2s;1-x^2)].\end{aligned}$$

Therefore,

$$\begin{aligned}S_{II} =& \left[\frac{1}{s-1}\frac{4}{x^2-1}+\frac{1}{s(2-s)}\frac{2(x^2+1)}{x^2}\right]F(2-s,1-s;3-2s;1-x^2)\\&+\left[\frac{1}{(1-s)(2-s)}\frac{4}{x^2-1}+\frac{1}{s(2-s)}\frac{4(x^2+1)}{x^2(x^2-1)}\right]F(1-s,1-s;2-2s;1-x^2)\\&+\left[\frac{1}{s(1-s)}\frac{2}{x^2(x^2-1)}-\frac{1}{s^2}\frac{2}{x^2-1}\right]F(2-s,-s;3-2s;1-x^2)\\&+\frac{(2-s)}{s^2(1-s)}\frac{2}{x^2(x^2-1)}F(-s,-s;3-2s;1-x^2)\\&+\left[-\frac{1}{s^2(1-s)}\frac{4}{x^2(x^2-1)}+\frac{1}{s^2}\frac{2}{x^2-1}\right]F(1-s,-s;3-2s;1-x^2).\end{aligned}$$

It is known that (see [Er], p.103, (28))

$$(c-a)F(a-1)+(2a-c-az+bz)F+a(z-1)F(a+1)=0.$$

Set $a=1-s$, $b=1-s$, $c=3-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(2-s,1-s;3-2s;1-x^2)\\ =&\frac{2-s}{1-s}\frac{1}{x^2}F(1-s,-s;3-2s;1-x^2)-\frac{1}{1-s}\frac{1}{x^2}F(1-s,1-s;3-2s;1-x^2).\end{aligned}$$

Set $a=1-s$, $b=-s$, $c=3-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(2-s,-s;3-2s;1-x^2)\\ =&\frac{2-s}{1-s}\frac{1}{x^2}F(-s,-s;3-2s;1-x^2)+\frac{1}{1-s}\frac{x^2-2}{x^2}F(1-s,-s;3-2s;1-x^2).\end{aligned}$$

Note that

$$\begin{aligned}&\frac{1}{(1-s)(2-s)}\frac{4}{x^2-1}+\frac{1}{s(2-s)}\frac{4(x^2+1)}{x^2(x^2-1)}\\ =&\frac{1}{s(1-s)(2-s)}\frac{4}{x^2-1}+\frac{1}{s(2-s)}\frac{4}{x^2(x^2-1)},\end{aligned}$$

and

$$-\frac{1}{s^2(1-s)}\frac{4}{x^2(x^2-1)}+\frac{1}{s^2}\frac{2}{x^2-1}-\frac{1}{s^2(1-s)}\frac{2(x^2-2)}{x^2(x^2-1)}=\frac{-1}{s(1-s)}\frac{2}{x^2-1}.$$

We have

$$\begin{aligned}S_{II}=&[-\frac{1}{s(1-s)}\frac{2}{x^2-1}-\frac{(2-s)}{(1-s)^2}\frac{4}{x^2(x^2-1)}+\frac{1}{s(1-s)}\frac{2(x^2+1)}{x^4}\\ &+\frac{1}{s(1-s)^2}\frac{2(x^2-2)}{x^4(x^2-1)}]F(1-s,-s;3-2s;1-x^2)\\ &+[\frac{1}{(1-s)^2}\frac{4}{x^2(x^2-1)}-\frac{1}{s(1-s)(2-s)}\frac{2(x^2+1)}{x^4}]F(1-s,1-s;3-2s;1-x^2)\\ &+[\frac{1}{s(2-s)}\frac{4}{x^2(x^2-1)}+\frac{1}{s(1-s)(2-s)}\frac{4}{x^2-1}]F(1-s,1-s;2-2s;1-x^2)\\ &+\frac{(2-s)}{s(1-s)^2}\frac{2}{x^4(x^2-1)}F(-s,-s;3-2s;1-x^2).\end{aligned}$$

It is known that (see [Er], p.103, (38))

$$c(1-z)F-cF(a-1)+(c-b)zF(c+1)=0.$$

Set $a = 1 - s$, $b = 1 - s$, $c = 2 - 2s$ and $z = 1 - x^2$, then

$$\begin{aligned}&F(1-s, 1-s; 2-2s; 1-x^2)\\=&\frac{1}{x^2}F(-s, 1-s; 2-2s; 1-x^2) + \frac{x^2-1}{2x^2}F(1-s, 1-s; 3-2s; 1-x^2).\end{aligned}$$

Note that

$$\begin{aligned}&\frac{1}{(1-s)^2}\frac{4}{x^2(x^2-1)} - \frac{1}{s(1-s)(2-s)}\frac{2(x^2+1)}{x^4} + \frac{1}{s(2-s)}\frac{2}{x^4} + \frac{1}{s(1-s)(2-s)}\frac{2}{x^2}\\=&\frac{1}{(1-s)^2}\frac{4}{x^2(x^2-1)} - \frac{1}{(1-s)(2-s)}\frac{2}{x^4}.\end{aligned}$$

Thus,

$$\begin{aligned}S_{II} =&[-\frac{1}{s(1-s)}\frac{2}{x^2-1} - \frac{(2-s)}{(1-s)^2}\frac{4}{x^2(x^2-1)} + \frac{1}{s(1-s)}\frac{2(x^2+1)}{x^4}\\&+\frac{1}{s(1-s)^2}\frac{2(x^2-2)}{x^4(x^2-1)}]F(1-s, -s; 3-2s; 1-x^2)\\&+[\frac{1}{(1-s)^2}\frac{4}{x^2(x^2-1)} - \frac{1}{(1-s)(2-s)}\frac{2}{x^4}]F(1-s, 1-s; 3-2s; 1-x^2)\\&+[\frac{1}{s(2-s)}\frac{4}{x^4(x^2-1)} + \frac{1}{s(1-s)(2-s)}\frac{4}{x^2(x^2-1)}]F(-s, 1-s; 2-2s; 1-x^2)\\&+\frac{(2-s)}{s(1-s)^2}\frac{2}{x^4(x^2-1)}F(-s, -s; 3-2s; 1-x^2).\end{aligned}$$

We know that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = -s$, $b = 1 - s$, $c = 3 - 2s$ and $z = 1 - x^2$, then

$$\begin{aligned}&F(-s, 1-s; 2-2s; 1-x^2)\\=&\frac{2-s}{2(1-s)}F(-s, 1-s; 3-2s; 1-x^2) - \frac{s}{2(1-s)}F(1-s, 1-s; 3-2s; 1-x^2).\end{aligned}$$

Note that

$$\begin{aligned}&\frac{1}{(1-s)^2}\frac{4}{x^2(x^2-1)} - \frac{1}{(1-s)(2-s)}\frac{2}{x^4} - \frac{1}{(1-s)(2-s)}\frac{2}{x^4(x^2-1)}\\&-\frac{1}{(1-s)^2(2-s)}\frac{2}{x^2(x^2-1)} = \frac{1}{(1-s)^2}\frac{2}{x^2(x^2-1)}.\end{aligned}$$

Therefore,

$$\begin{aligned}S_{II} =&[-\frac{1}{s(1-s)}\frac{2}{x^2-1}-\frac{(2-s)}{(1-s)^2}\frac{4}{x^2(x^2-1)}+\frac{1}{s(1-s)}\frac{2(x^2+1)}{x^4}\\&+\frac{1}{s(1-s)^2}\frac{2(x^2-2)}{x^4(x^2-1)}+\frac{1}{s(1-s)}\frac{2}{x^4(x^2-1)}+\frac{1}{s(1-s)^2}\frac{2}{x^2(x^2-1)}]\\&\times F(1-s,-s;3-2s;1-x^2)\\&+\frac{1}{(1-s)^2}\frac{2}{x^2(x^2-1)}F(1-s,1-s;3-2s;1-x^2)\\&+\frac{(2-s)}{s(1-s)^2}\frac{2}{x^4(x^2-1)}F(-s,-s;3-2s;1-x^2)\end{aligned}$$

Note that

$$-\frac{1}{s(1-s)}\frac{2}{x^2-1}+\frac{1}{s(1-s)}\frac{2}{x^4(x^2-1)}+\frac{1}{s(1-s)}\frac{2(x^2+1)}{x^4}=0,$$

and

$$\begin{aligned}&\frac{1}{s(1-s)^2}\frac{2(x^2-2)}{x^4(x^2-1)}+\frac{1}{s(1-s)^2}\frac{2}{x^2(x^2-1)}\\=&\frac{-1}{s(1-s)^2}\frac{4}{x^4(x^2-1)}+\frac{1}{s(1-s)^2}\frac{4}{x^2(x^2-1)}.\end{aligned}$$

We have

$$\begin{aligned}S_{II}=&\left[-\frac{(2-s)}{(1-s)^2}\frac{4}{x^2(x^2-1)}-\frac{1}{s(1-s)^2}\frac{4}{x^4(x^2-1)}+\frac{1}{s(1-s)^2}\frac{4}{x^2(x^2-1)}\right]\\&\times F(1-s,-s;3-2s;1-x^2)\\&+\frac{1}{(1-s)^2}\frac{2}{x^2(x^2-1)}F(1-s,1-s;3-2s;1-x^2)\\&+\frac{2-s}{s(1-s)^2}\frac{2}{x^4(x^2-1)}F(-s,-s;3-2s;1-x^2).\end{aligned}$$

In the formula (see [Er], p.103, (36))

$$(c-a-b)F-(c-a)F(a-1)+b(1-z)F(b+1)=0,$$

set $a=1-s$, $b=-s$, $c=3-2s$ and $z=1-x^2$, then

$$\begin{aligned}&F(1-s,1-s;3-2s;1-x^2)+\frac{2-s}{s}\frac{1}{x^2}F(-s,-s;3-2s;1-x^2)\\=&\frac{1}{s}\frac{2}{x^2}F(1-s,-s;3-2s;1-x^2).\end{aligned}$$

Now, we have

$$\begin{aligned}
&\frac{1}{(1-s)^2}\frac{2}{x^2(x^2-1)}F(1-s,1-s;3-2s;1-x^2)\\
&+\frac{(2-s)}{s(1-s)^2}\frac{2}{x^4(x^2-1)}F(-s,-s;3-2s;1-x^2)\\
=&\frac{1}{(1-s)^2}\frac{2}{x^2(x^2-1)}\frac{1}{s}\frac{2}{x^2}F(1-s,-s;3-2s;1-x^2)\\
=&\frac{1}{s(1-s)^2}\frac{4}{x^4(x^2-1)}F(1-s,-s;3-2s;1-x^2).
\end{aligned}$$

Hence,

$$\begin{aligned}
S_{II}=&[-\frac{(2-s)}{(1-s)^2}\frac{4}{x^2(x^2-1)}-\frac{1}{s(1-s)^2}\frac{4}{x^4(x^2-1)}+\frac{1}{s(1-s)^2}\frac{4}{x^2(x^2-1)}\\
&+\frac{1}{s(1-s)^2}\frac{4}{x^4(x^2-1)}]F(1-s,-s;3-2s;1-x^2)\\
=&\frac{1}{s}\frac{4}{x^2(x^2-1)}F(1-s,-s;3-2s;1-x^2)\\
=&\frac{4}{s}\frac{u^2}{u+1}F(1-s,-s;3-2s;-u^{-1}).
\end{aligned}$$

Therefore, we have

$$\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}(4u)^{s-3}S_{II}=\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}4^{s-2}u^{s-1}(u+1)^{-1}F(1-s,-s;3-2s;-u^{-1}).$$

Consequently,

$$\phi(Z,Z';k,s)=K(Z,Z';k,s)+K(Z,Z';k,2-s),$$

where $k=\pm1$.

□

Corollary 4.12. *The functional equation of Eisenstein series of weight k for the trivial group on $\mathfrak{S}_2$: If $Re(s)>1$, then the following identity holds:*

$$\begin{aligned}
&\int_{\partial\mathfrak{S}_2}P_k(Z,W;s)\rho(W,W')^{s+k-2}\rho(\overline{W},\overline{W'})^{s-k-2}dm(W)\\
=&\frac{\pi^{\frac{3}{2}}\Gamma(s-\frac{1}{2})}{(|k|+s-1)\Gamma(s)}4^{s-1}P_k(Z,W';2-s),
\end{aligned}\tag{4.31}$$

for $k=0,\pm1$, where $W'\in\partial\mathfrak{S}_2$.

Proof. The integral on the left-hand side is absolutely convergent. We set

$$P_{-k}(Z',W;2-s)=\left[\frac{\rho(Z',W)}{\rho(\overline{Z'},\overline{W})}\right]^{-k}\frac{\rho(Z')^{2-s}}{|\rho(Z',W)|^{2(2-s)}}$$

in Theorem 4.10. Next, we multiply both sides of the formula by $\rho(Z')^{s-2}$ and take the limit as $Z'\to W'\in\partial\mathfrak{S}_2$. This completes the proof.

□

We define the S-matrix of weight k as follows:

$$S(W,W';k,s):=\sum_{\gamma\in\Gamma}j(\gamma,W)^{k-s}j(\overline{\gamma},\overline{W})^{-k-s}\rho(\gamma(W),W')^{-k-s}\rho(\overline{\gamma(W)},\overline{W'})^{k-s}\tag{4.32}$$

for $W,W'\in\Omega(\Gamma)$ and $\mathrm{Re}(s)>\delta(\Gamma)$.

THEOREM 4.13. *Assume that Γ is convex cocompact and $\delta(\Gamma)<1$. Then the following functional equation for Eisenstein series of weight k holds:*

$$\begin{aligned}&\int_{\Gamma\backslash\Omega(\Gamma)}E(Z,W;k,s)S(W,W';-k,2-s)dm(W)\\&=\frac{\pi^{\frac{3}{2}}\Gamma(s-\frac{1}{2})}{(|k|+s-1)\Gamma(s)}4^{s-1}E(Z,W';k,2-s),\end{aligned}\tag{4.33}$$

for $1<Re(s)<2-\delta(\Gamma)$ and $k=0,\pm1$.

Proof. By Corollary 4.12,

$$\begin{aligned}&\int_{\partial\mathfrak{S}_2}E(Z,W;k,s)\rho(W,W')^{s+k-2}\rho(\overline{W},\overline{W'})^{s-k-2}dm(W)\\&=\frac{\pi^{\frac{3}{2}}\Gamma(s-\frac{1}{2})}{(|k|+s-1)\Gamma(s)}4^{s-1}E(Z,W';k,2-s).\end{aligned}\tag{4.34}$$

On the other hand,

$$\begin{aligned}\int_{\partial\mathfrak{S}_2}f(W)dW&=\int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma}f(\gamma(W))dm(\gamma(W))\\&=\int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma}|J(\gamma(W))|f(\gamma(W))dm(W).\end{aligned}$$

By Lemma 4.9, (3), the left hand side of (4.34) is equal to

$$\begin{aligned}&\int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma}|J(\gamma(W))|E(Z,\gamma(W);k,s)\rho(\gamma(W),W')^{s+k-2}\rho(\overline{\gamma(W)},\overline{W'})^{s-k-2}dm(W)\\ =&\int_{\Gamma\backslash\Omega(\Gamma)}\sum_{\gamma\in\Gamma}|j(\gamma,W)|^{-4}j(\gamma,W)^{s-k}j(\overline{\gamma},\overline{W})^{s+k}E(Z,W;k,s)\\ &\times\rho(\gamma(W),W')^{s+k-2}\rho(\overline{\gamma(W)},\overline{W'})^{s-k-2}dm(W)\\ =&\int_{\Gamma\backslash\Omega(\Gamma)}E(Z,W;k,s)S(W,W';-k,2-s)dm(W).\end{aligned}$$

□

By Theorem 4.10, we get the main theorem of this paper.

THEOREM 4.14. *Assume that* Γ *is convex cocompact and* $\delta(\Gamma)<1$. *Then the following product formula holds:*

$$\begin{aligned}&\int_{\Gamma\backslash\Omega(\Gamma)}E(Z,W;k,s)E(Z',W;-k,2-s)dm(W)\\ =&G(Z,Z';k,s)+G(Z,Z';k,2-s)\\ =&G(Z',Z;-k,s)+G(Z',Z;-k,2-s),\end{aligned}\tag{4.35}$$

for $\delta(\Gamma)<Re(s)<2-\delta(\Gamma)$ *and* $k=0$ *or* $k=\pm1$.

Appendix. Product formulas on $SL(2,\mathbb{R})$

In this appendix, by the same method as in the above argument we give the product formulas on $SL(2,\mathbb{R})$. For simplicity, we omit some details.

Let

$$\Delta_k=-(z-\overline{z})^2\frac{\partial^2}{\partial z\partial\overline{z}}-k(z-\overline{z})(\frac{\partial}{\partial z}+\frac{\partial}{\partial\overline{z}}).$$

Set $H_k(z,w)=(\overline{z}-w)^k(z-\overline{w})^{-k}$ and $P(z,\zeta)=\frac{\mathrm{Im}(z)}{|\overline{z}-\zeta|^2}$, where $z,w\in\mathbb{H}$ and $\zeta\in\partial\mathbb{H}=\mathbb{R}$. The Poisson kernel of weight k is defined as follows:

$$P_k(z,\zeta;s)=(\overline{z}-\zeta)^k(z-\zeta)^{-k}P(z,\zeta)^s.$$

$$E(z,\zeta;k,s):=\sum_{\gamma\in\Gamma}P_k(\gamma(z),\zeta;k,s),\quad \text{for}\quad \mathrm{Re}(s)>\delta(\Gamma).$$

Here Γ is a discrete subgroup of $SL(2,\mathbb{R})$. The point-pair invariant

$$u(z,z'):=\frac{|z-z'|^2}{4\mathrm{Im}(z)\mathrm{Im}(z')},\quad \sigma(z,z'):=\frac{|\overline{z}-z'|^2}{4\mathrm{Im}(z)\mathrm{Im}(z')}.$$

In fact, $u = \sigma - 1$.

If $f = H_k(z,w)\Phi(\sigma(z,w))$, then

$$\Delta_k f = H_k(z,w)\left[\sigma(\sigma-1)\Phi''(\sigma) + (2\sigma-1)\Phi'(\sigma) + \frac{k^2}{\sigma}\Phi(\sigma)\right].$$

By $\Delta_k f = s(s-1)f$, we have

$$\sigma(\sigma-1)\Phi''(\sigma) + (2\sigma-1)\Phi'(\sigma) + \left[\frac{k^2}{\sigma} - s(s-1)\right]\Phi(\sigma) = 0.$$

A solution is

$$\begin{aligned}\Phi(\sigma) =& P\left\{\begin{matrix} 0 & 1 & \infty & \\ -|k| & 0 & s & ;\sigma \\ |k| & 0 & 1-s & \end{matrix}\right\} \\ =& \sigma^{-|k|}(1-\sigma)^{|k|-s}F(s-|k|, s-|k|; 2s; -\frac{1}{\sigma-1}).\end{aligned}$$

Set

$$\begin{aligned}K(z,z';k,s) =& \frac{\sqrt{\pi}\Gamma(\frac{1}{2}-s)}{\Gamma(1-s)}4^{-s}\frac{s}{s-|k|}H_k(z,z') \\ & \times u^{|k|-s}(1+u)^{-|k|}F(s-|k|, s-|k|; 2s; -u^{-1}),\end{aligned}$$

where $u = u(z,z')$. The automorphic Green function of weight k is given by

$$G(z,z';k,s) = \sum_{\gamma\in\Gamma} K(z,\gamma(z');k,s), \quad \text{for} \quad \mathrm{Re}(s) > \delta(\Gamma).$$

Let

$$\phi(z,z';k,s) = \int_{\mathbb{R}} P_k(z,\zeta;s)P_{-k}(z',\zeta;1-s)d\zeta.$$

Then ϕ is a point-pair invariant covariant with respect to the weight k.

Put $z = iy$, $z' = iy'$, $y > 0$, $y' > 0$. Then

$$\begin{aligned}&\phi(iy,iy';k,s) \\ =& y^s {y'}^{1-s}\int_{-\infty}^{\infty}\left(\frac{\zeta+iy}{\zeta-iy}\right)^k\left(\frac{\zeta+iy'}{\zeta-iy'}\right)^{-k}|\zeta-iy|^{-2s}|\zeta-iy'|^{-2(1-s)}d\zeta \\ =& y^s {y'}^{1-s}\int_{-\infty}^{\infty}\exp(2ki\arctan\frac{y}{\zeta} - 2ki\arctan\frac{y'}{\zeta})\frac{1}{(\zeta^2+y^2)^s({y'}^2+\zeta^2)^{1-s}}d\zeta.\end{aligned}$$

By $\tan(\arctan\frac{y}{\zeta}-\arctan\frac{y'}{\zeta})=\frac{\zeta(y-y')}{\zeta^2+yy'}$,

$$\cos(2\arctan\left[\frac{(y-y')\zeta}{yy'+\zeta^2}\right])=\frac{(yy'+\zeta^2)^2-(y-y')^2\zeta^2}{(yy'+\zeta^2)^2+(y-y')^2\zeta^2},$$

and

$$\begin{aligned}&\cos(2k\arctan\left[\frac{(y-y')\zeta}{yy'+\zeta^2}\right])\\ =&\cos(k\arccos\frac{(yy'+\zeta^2)^2-(y-y')^2\zeta^2}{(yy'+\zeta^2)^2+(y-y')^2\zeta^2})\\ =&F(-k,k;\frac{1}{2};\frac{1}{2}\left[1-\frac{(yy'+\zeta^2)^2-(y-y')^2\zeta^2}{(yy'+\zeta^2)^2+(y-y')^2\zeta^2}\right])\\ =&F(-k,k;\frac{1}{2};\frac{(y-y')^2\zeta^2}{(y^2+\zeta^2)({y'}^2+\zeta^2)}),\end{aligned}$$

we have

$$\begin{aligned}\phi(iy,iy';k,s)=&2y^s{y'}^{1-s}\sum_{n=0}^{|k|}\frac{(-k)_n(k)_n}{n!(\frac{1}{2})_n}(y-y')^{2n}\\ &\times\int_0^\infty\zeta^{2n}(\zeta^2+y^2)^{-s-n}(\zeta^2+{y'}^2)^{s-n-1}d\zeta.\end{aligned}$$

Set $l=\frac{y'}{y}$, then $u=\frac{(l-1)^2}{4l}$. By the same method as the above argument for $GU(2,1)$, we have

$$\begin{aligned}&\phi(iy,iy';k,s)\\ =&\sum_{n=0}^{|k|}\frac{(-k)_n(k)_n}{n!(\frac{1}{2})_n}\Gamma(n+\frac{1}{2})[\frac{\Gamma(\frac{1}{2}-s)}{\Gamma(n+1-s)}(4u)^{-s}F(n+s,s;2s;-u^{-1})\\ &+\frac{\Gamma(s-\frac{1}{2})}{\Gamma(n+s)}(4u)^{-(1-s)}F(n+1-s,1-s;2-2s;-u^{-1})].\end{aligned}$$

(1) $k=0$,

$$\phi(iy,iy';k,s)=\frac{\sqrt{\pi}\Gamma(\frac{1}{2}-s)}{\Gamma(1-s)}(4u)^{-s}F(s,s;2s;-u^{-1})+(s\mapsto 1-s).$$

(2) $k=\pm1$,

$$S_I:=\frac{\sqrt{\pi}\Gamma(\frac{1}{2}-s)}{\Gamma(1-s)}(4u)^{-s}[F(s,s;2s;-u^{-1})+\frac{1}{s-1}F(s+1,s;2s;-u^{-1})].$$

By [Er], p.103, (28),

$$(c-a)F(a-1)+(2a-c-az+bz)F+a(z-1)F(a+1)=0,$$

set $a=s$, $b=s$, $c=2s$ and $z=-u^{-1}$, then

$$F(s+1,s;2s;-u^{-1})=\frac{u}{u+1}F(s-1,s;2s;-u^{-1}).$$

By [Er], p.103, (36),

$$(c-a-b)F-(c-a)F(a-1)+b(1-z)F(b+1)=0,$$

set $a=s$, $b=s-1$, $c=2s$ and $z=-u^{-1}$, then

$$\begin{aligned}&\frac{1}{s-1}\frac{u}{u+1}F(s,s-1;2s;-u^{-1})+F(s,s;2s;-u^{-1})\\=&\frac{s}{s-1}\frac{u}{u+1}F(s-1,s-1;2s;-u^{-1}).\end{aligned}$$

Thus,

$$S_I=\frac{\sqrt{\pi}\Gamma(\frac{1}{2}-s)}{\Gamma(1-s)}(4u)^{-s}\frac{s}{s-1}\frac{u}{u+1}F(s-1,s-1;2s;-u^{-1}).$$

Consequently,

$$\phi(iy,iy';k,s)=\frac{\sqrt{\pi}\Gamma(\frac{1}{2}-s)}{\Gamma(1-s)}4^{-s}\frac{s}{s-1}u^{1-s}(u+1)^{-1}F(s-1,s-1;2s;-u^{-1})+(s\mapsto 1-s).$$

Thus, we have the following theorem:

THEOREM. *Assume that* Γ *is convex cocompact and* $\delta(\Gamma)<\frac{1}{2}$. *Then the following product formula on* $SL(2,\mathbb{R})$ *holds:*

$$\int_{\Gamma\backslash\Omega(\Gamma)}E(z,\zeta;k,s)E(z',\zeta;-k,1-s)dm(\zeta)=G(z,z';k,s)+G(z,z';k,1-s),$$

for $\delta(\Gamma)<Re(s)<1-\delta(\Gamma)$ *and* $k=0,\pm 1$.

DEPARTMENT OF MATHEMATICS, PEKING UNIVERSITY
BEIJING 100871, P. R. CHINA
E-mail address: yanglei@sxx0.math.pku.edu.cn

References

[Ap] B. Apanasov, Deformations and stability in complex hyperbolic geometry, MSRI preprint, No. 1997-111.

[BB] W. L. Baily and A. Borel, Compactification of arithmetic quotients of bounded symmetric domains, Ann. of Math. **84** (1966), 442-528.

[BO] U. Bunke and M. Olbrich, The spectrum of Kleinian manifolds, math.DG/9810146.

[Co] K. Corlette, Hausdorff dimensions of limit sets I, Invent. Math. **102** (1990), 521-542.

[CoI] K. Corlette and A. Iozzi, Limit sets of discrete groups of isometries of exotic hyperbolic spaces, Trans. AMS **351** (1999), 1507-1530.

[El] J. Elstrodt, Die Resolvente zum Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene, I, Math. Ann. **203** (1973), 295-330; II, Math. Z. **132** (1973), 99-134; III, Math. Ann. **208** (1974), 99-132.

[ElGM] J. Elstrodt, F. Grunewald and J. Mennicke, *Groups Acting on Hyperbolic Space, Harmonic Analysis and Number Theory*, Springer-Verlag, 1998.

[Er] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher Transcendental Functions*, Vol. I, McGraw-Hill Book Company, Inc., 1953.

[Fa] J. D. Fay, Fourier coefficients of the resolvent for a Fuchsian group, J. Reine Angew. Math. **294** (1977), 143-203.

[FHP] R. Froese, P. Hislop and P. Perry, The Laplace operator on hyperbolic three manifolds with cusps of non-maximal rank, Invent. Math. **106** (1991), 295-333.

[Gir] G. Giraud, Sur certaines fonctions automorphes de deux variables, Ann. Ec. Norm., **(3)**, **38** (1921), 43-164.

[Go] W. M. Goldman, *Complex Hyperbolic Geometry*, Oxford Mathematical Monographs, Oxford, 1999.

[H] D. Hejhal, *The Selberg Trace Formula for $PSL(2,\mathbb{R})$*, Vol. **1**, Lecture Notes in Math. **548**, Springer-Verlag, 1976; Vol. **2**, Lecture Notes in Math. **1001**, Springer-Verlag, 1983.

[He] J. C. Hemperly, The parabolic contribution to the number of linearly independent automorphic forms on a certain bounded domain, Amer. J. Math. **94** (1972), 1078-1100.

[HP] S. Hersonsky and F. Paulin, On the volumes of complex hyperbolic manifolds, Duke Math. J. **84** (1996), 719-737.

[Ho1] R. P. Holzapfel, *Geometry and Arithmetic Around Euler Partial Differential Equations*, Dt. Verl. d. Wiss., Berlin/Reidel, Dordrecht 1986.

[Ho2] R. P. Holzapfel, *The Ball and Some Hilbert Problems*, Lectures in Mathematics, ETH Zürich, Birkhäuser, 1995.

[K] M. Kashiwara, A. Kowata, K. Minemura, K. Okamoto, T. Oshima and M. Tanaka, Eigenfunctions of invariant differential operators on a symmetric space, Ann. of Math. 107(1978), 1-39.

[Ko] A. Korányi, The Poisson integral for generalized half-planes and bounded symmetric domains, Ann. of Math. **82** (1965), 332-350.

[KR] A. Korányi and H. M. Reimann, Quasiconformal mappings on the Heisenberg group, Invent. Math. **80** (1985), 309-338.

[KW] A. Korányi and J. A. Wolf, Realization of hermitian symmetric spaces as generalized half-planes, Ann. of Math. **81** (1965), 265-288.

[Ku] T. Kubota, *Elementary Theory of Eisenstein Series*. Wiley, New York, 1973.

[LR] R. P. Langlands and D. Ramakrishnan, *The Zeta Functions of Picard Modular Surfaces*, Les Publications CRM, Université de Montréal, 1992.

[M] H. Maass, Die Differentialgleichungen in der Theorie der elliptischen Modulfunktionen, Math. Ann. **125** (1953), 235-263.

[Ma1] N. Mandouvalos, Spectral theory and Eisenstein series for Kleinian groups, Proc. London Math. Soc. (3) **57** (1988), 209-238.

[Ma2] N. Mandouvalos, *Scattering Operator, Eisenstein Series, Inner Product Formula and "Maass-Selberg" Relations for Kleinian Groups*, Memoirs of A.M.S. **78**, No. **400** (1989).

[P] J. R. Parker, On the volumes of cusped, complex hyperbolic manifolds and orbifolds, Duke Math. J. **94** (1998), 433-464.

[Pa1] S. J. Patterson, The limit set of a Fuchsian group, Acta Math. **136** (1976), 241-273.

[Pa2] S. J. Patterson, The Laplacian operator on a Riemann surface, I, Comp. Math. **31** (1975), 83-107; II, Comp. Math. **32** (1976), 71-119, III, Comp. Math. **33** (1976), 227-259.

[Pa3] S. J. Patterson, The Selberg zeta function of a Kleinian group. In *Number Theory, Trace Formulas, and Discrete Groups*: Symposium in honor of Atle Selberg, Olso, Norway (1987), 409-442.

[Pe1] P. A. Perry, The Laplace operator on a hyperbolic manifold. II. Eisenstein series and the scattering matrix, J. Reine Angew. Math. **398** (1989), 67-91.

[Pe2] P. A. Perry, The Selberg zeta function and a local trace formula for Kleinian groups, J. Reine Angew. Math. **410** (1990), 116-152.

[Pic1] E. Picard, Sur une extension aux fonctions de deux variables du problème de Riemann relatif aux fonctions hypergeométriques séries hypergeométriques, Ann. Ec. Norm. **10** (1881), 305-322.

[Pic2] E. Picard, Sur les fonctions hyperfuchsiennes provenant des séries hypergeométriques de deux variables, Ann. Ec. Norm. **III 2** (1885), 357-384.

[Ro] W. Roelcke, Das Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene, I, Math. Ann. **167** (1966), 292-337; II, Math. Ann. **168** (1967), 261-324.

[Se] A. Selberg, Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series, J. Indian Math. Soc. B. **20** (1956), 47-87.

[Sh1] G. Shimura, The arithmetic of automorphic forms with respect to a unitary group, Ann. of Math. **107** (1978), 569-605.

[Sh2] G. Shimura, L-functions and eigenvalue problems. In *Algebraic Analysis, Geometry, and Number Theory*, (J. I. Igusa, ed.), Johns Hopkins University Press, Baltimore, 1989, 341-396.

[Su] D. Sullivan, The density at infinity of a discrete group of hyperbolic motions, Publ. Math. I.H.E.S. **50** (1979), 171-209.